



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





Chemical Library


Q II

271


.N357







THE IDENTIFICATION OF
ORGANIC COMPOUNDS



THE IDENTIFICATION OF ORGANIC COMPOUNDS

By

G. B. NEAVE, M.A., D.Sc. (St. Andrews)

and

I. M. HEILBRON, Ph.D. (Leipzig), F.I.C.

LECTURERS AND DEMONSTRATORS, DEPARTMENT
OF CHEMISTRY, GLASGOW AND WEST OF
SCOTLAND TECHNICAL COLLEGE



NEW YORK
D. VAN NOSTRAND COMPANY
23 MURRAY and 27 WARREN STREETS
1911

BUTLER & TANNER,
THE SELWOOD PRINTING WORKS,
FROME, AND LONDON.

Preface

IN teaching practical organic chemistry we have found the want of a convenient text-book dealing with the identification of simple organic compounds, such as is required by students working for the Intermediate and Final (Branch d) Examinations of the Institute of Chemistry. Moreover, many of the reactions and physical constants are not easily accessible, but are only to be obtained by a diligent and often tedious search through some of the larger books of reference.

In this small volume we have endeavoured to bring together in a convenient form the principal reactions and physical constants of the most important organic substances. Our aim has been to eliminate, as far as possible, guess-work on the part of the student, and to provide him with methods by which he can readily detect the more important groups in the compound, assign it to its class, and then complete its identification by referring to the section dealing with the class to which it belongs. Wherever

possible, an easily prepared derivative is described under each compound.

In many cases descriptions of operations are brief, as we assume that the student has already attended lectures on Organic Chemistry and worked through a satisfactory course of Preparations.

The scheme adopted in this book having given satisfactory results in the laboratories of this College, we now venture to give it wider publicity.

We are deeply indebted to Professor G. G. Henderson for the interest he has taken in the work, and for a number of valuable suggestions.

G. B. N.

I. M. H.

GLASGOW,
April, 1911.

Contents

SECTION	PAGE
I PRELIMINARY TESTS	1
II TESTS FOR THE ELEMENTS	4
III GROUP REACTIONS	8
IV HYDROCARBONS	14
V ALCOHOLS	19
VI ETHERS	22
VII PHENOLS	23
VIII ALDEHYDES	29
IX KETONES	36
X ACIDS	38
XI AROMATIC SULPHONIC ACIDS	48
XII ACID ANHYDRIDES	52
XIII ACID HALIDES	53
XIV ACID AMIDES	53
XV ACID IMIDES.	55
XVI ACID ANILIDES	55
XVII ESTERS.	56

SECTION	PAGE
XVIII QUINONES	58
XIX CARBOHYDRATES	60
XX GLUCOSIDES	61
XXI AMINES	62
XXII NITRO COMPOUNDS	68
XXIII NITROSO COMPOUNDS	72
XXIV NITRILES AND ISONITRILES	74
XXV ISOCYANATES.	75
XXVI UREAS AND UREIDES	75
XXVII URIC ACID GROUP	78
XXVIII HALOGEN COMPOUNDS	80
XXIX AZO COMPOUNDS	83
XXX PYRIDINE AND QUINOLINE GROUP	85
XXXI ALKALOIDS	86
XXXII SULPHUR COMPOUNDS	89
XXXIII TERPENES AND ALLIED COMPOUNDS	94
XXXIV ALBUMINS AND PROTEIDS	97
APPENDIX	99
INDEX	101

I. PRELIMINARY TESTS.

1. ORGANIC compounds are often characterized by their appearance and smell, and the experienced student may frequently be able to classify and in some cases to identify a substance by means of these physical properties.

Colour is generally induced in a compound by the presence of the following common groups :—

(a) Nitro group. The compound is then generally yellow.

(b) Nitroso group. The compound in the fused state or in solution (if monomolecular) is blue to green in colour.

(c) Azo group and other such related groups. The compound is generally highly coloured.

(d) Compounds having a quinonoid structure are usually strongly coloured. The common quinones are all deep yellow.

2. If the substance is a solid, a small quantity is heated on platinum foil and the changes which occur are carefully noted.

(a) A sooty flame indicates a high percentage of carbon in the substance. The compound is then probably one of the aromatic series.

(b) An incombustible residue indicates the presence of a metal or some inorganic matter in the compound, which is then tested for in the usual manner.

It must be remembered, however, that the metals Mercury and Arsenic, likewise Ammonium salts, are volatile.

3. The solubility of the substance in water is tested.

(a) With the exception of certain salts, substances containing hydroxyl radicles usually dissolve.

(b) The following classes of compounds are also generally soluble :—Lower alcohols, aldehydes, ketones, monobasic acids, polybasic acids, substituted acids, carbohydrates (except starch and cellulose), lower amines and amides, urea and its homologues, thioureas, cyanates, alkyl sulphates.

(c) The solution is tested with litmus.

(i) *An acid reaction* indicates the presence of a carboxyl or sulphonic group in the substance. A salt of a weak base would be hydrolytically dissociated in dilute solution with a resulting acid reaction.

Acid chlorides are rapidly, acid anhydrides slowly, decomposed by water, especially on warming.

(ii) *An alkaline reaction* usually indicates the presence of a free base.

4. A small quantity of the substance (about 0.25 gram) is heated with a large excess of soda-lime in a hard glass test-tube fitted with a cork and a short right-angled delivery tube.

(a) Many nitrogen compounds evolve ammonia. An amine is liberated from an amine salt.

(b) Formates are decomposed with evolution of hydrogen.

(c) Hydrocarbons are produced from carboxy acids or their salts.

(d) Phenols are formed from hydroxy aromatic acids or their salts.

(e) A smell of burnt sugar is observed from carbohydrates, glucosides and many higher acids such as citric acid, tartaric acid, malic acid, tannic acid and gallic acid.

5. The substance is warmed with concentrated caustic soda solution.

(a) Ammonia is evolved from ammonium salts, amides, and imides.

(b) Amines are liberated from their salts. The lower aliphatic amines have characteristic ammoniacal fishy odours and are inflammable. The aromatic amines are insoluble oils or solids.

(c) Acetyl derivatives of amines are decomposed, liberating the free amine. The decomposition is only accomplished by prolonged heating.

(d) Esters are slowly hydrolysed with liberation of the alcohol. The vapours of the lower alcohols are inflammable. The pleasant characteristic odour of the ester disappears.

(e) Acids, phenols and nitrophenols dissolve with formation of a salt. The nitrophenols give red or yellow solutions.

(f) Alkaloids are precipitated from solutions of

their salts. Morphine is only slowly precipitated from a solution of its salt and is readily soluble in excess of the reagent.

(g) Aliphatic aldehydes, with the exception of formaldehyde, are converted into resins.

(h) Solutions of glucose and lactose are turned brown.

6. Dilute sodium carbonate solution is added to a small quantity of the substance contained in a test-tube, and the mixture is gently warmed. Acids, chlorophenols, nitrophenols and polyhydric phenols readily dissolve. Monohydric phenols are insoluble.

7. In order to ascertain if the substance is a saturated or unsaturated compound, a small quantity is dissolved in chloroform or in some suitable solvent having no action on bromine and a dilute solution of bromine in the same solvent is added, drop by drop. If the bromine is instantly decolourized without a simultaneous evolution of hydrobromic acid, the presence of a double or treble bond is shown, the substance being unsaturated.

II. TESTS FOR THE ELEMENTS.

Oxygen cannot be detected by any direct test, and the indications of its presence in a substance must be inferred from the foregoing experiments.

It is not generally necessary to make any special tests for **Carbon** and **Hydrogen**.

The presence of **Nitrogen** in an organic substance is indicated by heating a small quantity of the compound in a hard glass test-tube with excess of soda-lime. If ammonia is evolved, the presence of nitrogen is proved.

As, however, nitro-compounds, azo-compounds and some other nitrogen compounds do not respond to this test, the following delicate test must be applied before the absence of nitrogen may be considered conclusively proved.

A small piece of sodium, about the size of a pea, is dropped into a dry hard glass test-tube and gently warmed till it melts, after which a little of the substance, about 0.1 gram, is introduced in such a manner that it falls directly on the sodium. If the substance is a liquid it is allowed to fall drop by drop on the sodium, care being taken to prevent it touching the sides of the tube. After the first violent action has ceased, the tube is heated in a Bunsen flame till all apparent change is over.

The tube is allowed to cool and 10 cc. water are carefully added. The mixture is boiled for a few minutes and, if necessary, filtered from any carbonaceous matter.


If the filtrate be divided into three portions the presence of a **halogen** or of **sulphur** in the substance may also be directly tested.

One portion of the original solution or filtrate is now treated with a few drops of caustic soda solu-


tion, a little freshly-prepared ferrous sulphate solution and finally a few drops of ferric chloride solution, and the whole well shaken. On carefully acidifying with concentrated hydrochloric acid a blue solution or precipitate of Prussian blue is formed, proving the presence of nitrogen in the original substance.

Note.—With substances containing only a small percentage of nitrogen a greenish-blue solution is frequently obtained which may only show the Prussian blue precipitate after standing for a considerable time.

The formation of the Prussian blue depends on the fact that the nitrogen and carbon combine with the sodium to yield sodium cyanide. This compound in presence of ferrous hydroxide combines to give sodium ferrocyanide—



After the addition of ferric chloride and acid the insoluble Prussian blue is precipitated—



After testing the one portion for the presence of nitrogen another portion is acidified with nitric acid, the solution boiled to expel any hydrocyanic acid present, and then silver nitrate solution is added. If a precipitate is formed the presence of a halogen in the original substance is indicated and its nature may be determined by submitting the precipitate to the usual examination for the detection of chlorine, bromine, or iodine. This test depends on the

formation of sodium halide produced by combination of any halogen in the original substance with the metallic sodium.

Another simple test to prove the presence of a halogen in a substance is obtained by wrapping a small piece of copper oxide within a stout copper wire, heating it in a Bunsen flame until no green colouration is produced, and allowing it to cool. A very small quantity of the substance is then placed on the oxide, which is again heated in the flame, when, if a halogen be present, a green colouration is observed, due to the formation of a volatile halogen compound of copper. This test is not absolutely conclusive, as some non-halogen-compounds appear able to produce a similar colouration. Further, by this test, no information is obtained as to which halogen is present.

Sulphur in a compound is best detected by taking the third portion from the sodium extract as already described, and placing a drop of the alkaline solution on a silver coin. If the original substance contained sulphur, sodium sulphide will have been produced, forming on the silver coin a black precipitate of silver sulphide.

The presence of sodium sulphide in the alkaline filtrate is also readily detected by the addition of sodium nitroprusside solution, which produces a deep violet colouration with an alkali sulphide.

Phosphorus may be detected by fusing a small quantity of the substance with a mixture of equal portions of potassium nitrate and carbonate on a

piece of platinum foil. The residue is dissolved in water, acidified with a few drops of concentrated nitric acid and a phosphate tested for in the usual way with ammonium molybdate.

III. GROUP REACTIONS.

FROM the foregoing preliminary tests the student will have deduced the type of compound with which he is dealing, or at any rate something of its nature. The following tests for individual groups may now be performed :—

HYDROXYL GROUP.

(a) A small quantity of the compound is heated for about fifteen minutes with an excess of acetic anhydride.

If a hydroxyl group is present an acetyl derivative (oily or crystalline) is thus formed which may then be separated, washed with dilute sodium carbonate solution and finally hydrolysed by heating under a reflux condenser with caustic soda solution. The acetyl compound is, in this way, decomposed with formation of an acetate which can readily be recognized by applying the usual tests for acetic acid or acetates (*see* page 38).

(b) SCHOTTEN-BAUMANN REACTION.—About 1

gram of the compound is vigorously shaken up with about 1 cc. benzoyl chloride and sufficient dilute caustic soda solution to render the mixture alkaline. The product is poured into water and the crystalline benzoate separated by filtration, washed with water and then hydrolysed with caustic soda solution. After hydrolysis an alkali benzoate is formed, which can be detected in the usual manner (*see* page 41).

If required, the benzoyl derivative may be purified by recrystallization from alcohol, and its melting point determined.

METHOXY AND ETHOXY GROUPS.

Seven cc. hydriodic acid (Sp. Gr. 1.7) are added to about 0.2 gram of the compound in a test-tube fitted with cork and delivery tube, the end of which dips into an alcoholic solution of silver nitrate. The mixture is very gently heated in a glycerine bath to 140° and, if a methoxy or ethoxy group be present, methyl or ethyl iodide distils over and produces a precipitate of silver iodide in the alcoholic silver nitrate solution.

ALDEHYDE GROUP.

(a) TOLLEN'S REACTION.—Ammonium hydroxide solution is cautiously added to about 2 cc. silver nitrate solution until the precipitate which is first formed is just redissolved. One cc. caustic soda solution is next run in and then 2 or 3 drops of

the supposed aldehyde solution. If an aldehyde group is present an immediate deposit of metallic silver is produced in the cold.

(b) SCHIFF'S REACTION.—A few drops of the aldehyde are poured into about 5 cc. of a solution of Schiff's reagent (magenta solution which has been decolourized by sulphur dioxide). The red colouration of the magenta is immediately restored.

It should be noted that some ketones if present in large quantity react similarly, although the reappearance of the colour is more gradual and shows first after the mixture has stood for some time.

(c) Fehling's solution is rapidly decomposed with precipitation of red cuprous oxide on being warmed with an aliphatic aldehyde.

Aromatic aldehydes do not give this reaction.

(d) If about 1 cc. of an aliphatic aldehyde is warmed with 2 cc. concentrated caustic potash solution, a brown precipitate of aldehyde resin separates out.

Formaldehyde and aromatic aldehydes do not react in this manner, but yield mixtures of alcohols and acids, both containing the same number of carbon atoms as the original aldehyde.



(e) If 1 cc. of an aldehyde is vigorously shaken up in a test-tube with a few cc. of a cold saturated solution of sodium hydrogen sulphite¹ a precipitate

¹ See Appendix, page 100.

of the aldehyde "bisulphite compound" is formed. Sodium carbonate solution liberates the aldehyde from the acid sulphite compound.

Some ketones produce similar crystalline compounds.

KETONE GROUP.

If an aldehyde group is absent the presence of a ketone group may be confirmed by the preparation of the semicarbazone of the ketone (*see* page 30 for this preparation).

AMINO GROUP.

(a) CARBYLAMINE TEST : —About 0.1 gram of the substance is mixed with 3 drops of chloroform and 2 cc. alcoholic potash, and the whole carefully warmed. The characteristic disgusting carbylamine odour is produced if the substance contains an amino group.

(b) A small quantity of the compound is dissolved in a few cc. hydrochloric acid and sodium nitrite solution is then added until an excess of free nitrous acid is present. On warming, the amino group is converted into hydroxyl, with a vigorous evolution of nitrogen.

Aliphatic amines and aromatic amines having the amino group in a side chain produce alcohols.

Aromatic primary amines with the amino group in the nucleus form phenols.

To distinguish between the two classes of aromatic amines, the amine is diazotized (*see* page 64) in the cold, and a solution of β -naphthol in caustic soda added. An intensely coloured azo dye is produced if the amino group is in the nucleus.

IMINO GROUP.

If sodium nitrite is added to an acid solution of the compound, a yellow precipitate or solution of a nitrosoamine is formed, which may be confirmed by applying LIEBERMANN'S TEST for nitrosoamines :—

The precipitate or solution containing the nitrosoamine is extracted with ether, the ethereal solution washed and dried rapidly over calcium chloride and the ether then blown off. To the residue, oil or solid, is now added a small quantity of phenol and 4 to 5 drops concentrated sulphuric acid, when a deep greenish black colouration is produced. On pouring the mixture into water a red solution is formed, which is changed to blue or green on addition of alkalies.

Note.—With aliphatic tertiary amines no reaction results on addition of nitrous acid, but dialkylanilines interact readily, intensely coloured green nitroso compounds being formed, the NO-group displacing hydrogen of the benzene nucleus from the para position to the nitrogen atom. Substances of this type do not give Liebermann's reaction.

The hydrochlorides of these para-nitroso bodies,

which would be precipitated on addition of sodium nitrite to a hydrochloric acid solution of a dialkylamine, are generally yellow in colour and must not be confounded with the secondary nitrosoamines.

NITRO GROUP.

A small quantity of the substance is mixed in a test-tube with about 2 cc. concentrated hydrochloric acid and to this 1 gram zinc dust is gradually added in small quantities, and the mixture finally warmed. About 5 cc. water are now run in and then concentrated caustic alkali until the precipitate at first formed is redissolved. The amine produced is extracted with ether, and the presence of the amino group can be recognized by the tests given in the preceding paragraph.

NITRILES AND ISONITRILES.

(a) A small quantity of the substance is heated with 20 cc. concentrated hydrochloric acid under a reflux condenser. The reaction mixture is then made alkaline with caustic soda and heated.

Ammonia is evolved from nitriles, whereas the isonitriles yield under the same conditions sodium formate and an amine.

(b) MENDIUS' REACTION FOR NITRILES.¹—Nitriles, when reduced in a similar manner to that described for the reduction of nitro-compounds, yield amines which can readily be isolated and identified.

¹ See page 74.

AZO-GROUP.

Azo dye-stuffs when vigorously reduced, yield amines, either simple or complex. About 1 gram of the substance is treated with about 20 cc. of a concentrated solution of stannous chloride in hydrochloric acid. Caustic soda solution is then added until the precipitate first formed is just redissolved. Free amines are liberated which may be separated, and the usual tests for the amino group applied.

IV. HYDROCARBONS.

THESE are generally colourless liquids or solids, practically insoluble in water, soluble in alcohol and ether. The solubility in the last mentioned solvents decreases with increase in the molecular weight of the compound.

ALIPHATIC HYDROCARBONS.

n-Pentane, C_5H_{12} , is an exceptionally stable colourless liquid, boiling at 36–37°.

n-Hexane, C_6H_{14} , boils at 69–70°.

n-Heptane, C_7H_{16} , boils at 97–98°.

n-Octane, C_8H_{18} , boils at 124°.


Isoprene, $\begin{smallmatrix} CH_2 \\ CH_2 \end{smallmatrix} > C-CH=CH_2$, is a colourless liquid

boiling at 37° , obtained by dry distillation of caoutchouc. Concentrated hydrochloric acid converts isoprene into a substance very similar to, if not identical with, caoutchouc.

AROMATIC HYDROCARBONS.

Benzene, C_6H_6 , B.P. $80-81^{\circ}$, M.P. 5.4° . The pure hydrocarbon has a peculiar, not unpleasant smell. It is exceedingly stable towards oxidizing agents.

If 1 gram of the hydrocarbon be added to a mixture of 3 grams concentrated sulphuric acid and 2 grams concentrated nitric acid (Sp. Gr. 1.4), and the mixture warmed on a water-bath for about $\frac{1}{2}$ hour at 60° , nitro-benzene is obtained, and on cooling may be separated from the lower layer of acids, washed, dried and identified [(see page 70)



Toluene, $C_6H_5 \cdot CH_3$, B.P. 110° . Toluene on treatment with chromic acid yields benzoic acid.

About 1 gram of the substance is heated under a reflux condenser with a mixture consisting of 2 parts potassium dichromate, 3 parts concentrated sulphuric acid and 3 of water. After oxidation the solution is filtered while hot, when on cooling benzoic acid separates out and may be tested for as shown on page 41.

***o*-Xylene**, $C_6H_4 \begin{smallmatrix} \diagup CH_3 \\ \diagdown CH_3 \end{smallmatrix}$, boils at 142° . With dilute potassium permanganate it yields phthalic acid.

A small quantity of the hydrocarbon is heated under a reflux condenser for some time with dilute alkaline potassium permanganate. The solution is then, if necessary, treated with sulphurous acid to remove excess of potassium permanganate, filtered off from the separated manganese dioxide and acidified. The acid is then extracted with ether, dried with calcium chloride and the tests for phthalic acid applied (*see* page 42).

m-Xylene and *p*-Xylene, boiling points 139° and 138° respectively, are distinguished from each other by the fact that the meta-compound is sulphonated in the cold with sulphuric acid, while the para-compound remains unchanged.

Ethylbenzene, $C_6H_5 \cdot C_2H_5$, boils at 134° . Chromic acid oxidizes it to benzoic acid.

Mesitylene, $C_6H_3(CH_3)_3$ 1 : 3 : 5, is an agreeably smelling liquid boiling at $163-164^{\circ}$. With chromic acid mesitylene is decomposed into acetic acid.

Cumene, $C_6H_5 \cdot CH(CH_3)_2$, is a colourless liquid boiling at $152-153^{\circ}$. On oxidation with chromic acid or dilute nitric acid it is converted into benzoic acid.

Cymene, $C_6H_4 \begin{matrix} \swarrow CH_3 \\ \searrow CH(CH_3)_2 \end{matrix} \begin{matrix} (1) \\ (4) \end{matrix}$, is a pleasant smelling liquid boiling at 175° .

Diphenyl, $C_6H_5 \cdot C_6H_5$, is a colourless, crystalline solid, melting at 71° , B.P. 254° ; when oxidized with chromic acid it is converted into benzoic acid.

Diphenylmethane, $C_6H_5 \cdot CH_2 \cdot C_6H_5$, is a crystalline solid, M.P. 26.5° . On oxidation with chromic acid

it is converted into benzophenone which may be extracted with ether and identified (page 37).

Stilbene, $C_6H_5 \cdot CH : CH \cdot C_6H_5$, crystallizes in colourless needles, M.P. $124-125^\circ$. It is unsaturated and readily adds on two atoms of bromine forming stilbene dibromide, $C_6H_5 \cdot CHBr \cdot CHBr \cdot C_6H_5$, M.P. 137° . Oxidation with chromic acid or alkaline permanganate yields benzoic acid.

Styrene, $C_6H_5 \cdot CH : CH_2$, is a colourless liquid which boils at 145° . It is oxidized to benzoic acid with chromic acid mixture.

Phenylacetylene, $C_6H_5 \cdot C : CH$, is a colourless liquid boiling at $139-140^\circ$. It yields a yellow copper compound when mixed with ammoniacal cuprous chloride solution. On oxidation with chromic acid, benzoic acid is obtained.

Naphthalene, $C_{10}H_8$, crystallizes in lustrous leaflets which melt at 80° and boil at 218° . It has a characteristic smell, and is very volatile. On addition of a solution of the hydrocarbon in benzene to a solution of picric acid in the same solvent, a yellow crystalline compound, naphthalene picrate, $C_{10}H_8 \cdot C_6H_2(NO_2)_3OH$ melting at 149° separates out on standing. Naphthalene on boiling with dilute nitric acid is oxidized, yielding phthalic acid and carbon dioxide.


Anthracene, $C_{14}H_{10}$, crystallizes in almost colourless lustrous plates, possessing a beautiful violet fluorescence. It melts at 213° , boils at 360° . With picric acid it forms the molecular compound $C_{14}H_{10} \cdot C_6H_2(NO_2)_3OH$ which is deposited from

benzene solution in ruby-red needles melting at 138° .

Anthracene is readily oxidized to anthraquinone by dissolving the hydrocarbon in warm glacial acetic acid and adding to the solution about double the weight of chromic anhydride, and then boiling for a short time. The mixture is largely diluted with water, the anthraquinone filtered off, washed with dilute sulphuric acid, then with water and finally dried. M.P. 273° .

Phenanthrene, $C_{14}H_{10}$, forms colourless leaflets melting at 99° . It is readily soluble in alcohol, while the isomeric anthracene is only sparingly soluble.

It combines with picric acid to give a yellow compound, phenanthrene picrate, melting at 144° . With chromic anhydride it yields phenanthraquinone, M.P. 205° .



leaflets with a violet fluorescence. It melts at 113° . The picrate obtained by mixing benzene solutions of fluorene and picric acid melts at $79-80^{\circ}$.

V. ALCOHOLS.

THE common monohydric alcohols are, as a rule, colourless liquids (diphenyl and triphenyl carbinols are solids) with a neutral reaction and possessing a characteristic smell and taste. The lower members are soluble in water, but the solubility rapidly decreases with increase of molecular weight. The Sp. Gr. is always less than that of water.

The polyhydric alcohols are oily liquids or crystalline solids, all readily soluble in water, sparingly soluble or insoluble in ether.

PRIMARY MONOHYDRIC ALCOHOLS.

Methyl alcohol, CH_3OH , is a colourless, mobile liquid boiling at 66.5° . It burns with a bluish non-luminous flame. If a mixture of methyl alcohol with dilute sulphuric acid and potassium dichromate be distilled, the distillate contains formic acid which may be identified (page 38).

When methyl alcohol is heated with salicylic acid and concentrated sulphuric acid, methyl salicylate is formed and can be identified by its odour. On warming methyl alcohol with paranitrobenzoyl chloride, the methyl ester is formed, melting at 96° .


Ethyl Alcohol, $\text{C}_2\text{H}_5\text{OH}$, boils at 78° .

1. When ethyl alcohol is mixed with a little dilute sulphuric acid and potassium dichromate and the mixture warmed, the alcohol is oxidized to acetalde-

hyde which can be recognized by its peculiar choking smell.

2. The formation of Iodoform serves as a delicate test for ethyl alcohol.¹ To a solution of alcohol a little sodium carbonate is added and the mixture warmed to about 60°. To this is now added drop by drop, a strong solution of iodine in potassium iodide until, after shaking, the liquid remains faintly brown. Iodoform separates out, having a characteristic odour and melting at 119°.

3. The para-nitrobenzoic-ester,



is readily obtained as a crystalline solid melting at 57°.

Propyl alcohol, $\text{C}_3\text{H}_7\text{OH}$, boils at 97·4°. It is miscible in all proportions with water, but on addition of calcium chloride and other easily soluble salts it separates out from its aqueous solution.

Normal Butyl Alcohol, $\text{C}_4\text{H}_9\text{OH}$, is a liquid with an agreeable odour boiling at 117°.

Isobutyl Alcohol, $\begin{matrix} \text{CH}_3 \\ \text{CH}_3 \end{matrix} > \text{CH}\cdot\text{CH}_2\text{OH}$, is a liquid possessing a fusel-oil odour. B.P. 108°.

Normal Amyl Alcohol, $\text{C}_5\text{H}_{11}\text{OH}$, is a liquid almost insoluble in water, boiling at 137°.

Allyl Alcohol, $\text{C}_3\text{H}_5\text{OH}$ ($\text{CH}_2 : \text{CH}\cdot\text{CH}_2\text{OH}$), is a mobile liquid with a pungent odour, boiling at 96·5°. It has the properties not only of a primary alcohol, but also of an unsaturated compound.

¹ Acetone also gives the Iodoform test.

On distilling allyl alcohol with chromic acid, formic acid (page 38) distils over.

Benzyl Alcohol, $C_6H_5 \cdot CH_2OH$, is a colourless liquid, with a faint aromatic odour and boils at 206° . It is sparingly soluble in water, readily in alcohol and ether. On oxidation with dilute nitric acid it yields first benzaldehyde and then benzoic acid.

SECONDARY MONOHYDRIC ALCOHOLS.

Isopropyl Alcohol ($CH_3 \cdot CHOH \cdot CH_3$), boils at $82-83^\circ$. Oxidizing agents convert it into acetone.

Diphenyl Carbinol, benzhydrol, $(C_6H_5)_2CHOH$, crystallizes in silky needles, very sparingly soluble in water, M.P. $67-68^\circ$. The acetate melts at $41-42^\circ$.

TERTIARY MONOHYDRIC ALCOHOLS.

Tertiary Butyl Alcohol, $(CH_3)_3COH$, is a colourless solid melting at 25° , boiling at $83-84^\circ$.

Triphenyl Carbinol, $(C_6H_5)_3COH$, M.P. 159° . The acetate melts at 99° .

POLYHYDRIC ALCOHOLS.

Ethylene Glycol, $CH_2OH \cdot CH_2OH$, is a thick colourless liquid boiling at $197-198^\circ$. When glycol is heated with solid caustic potash to 250° , hydrogen is evolved and potassium oxalate formed which may be identified (*see* page 39).

Glycerol, $C_3H_5(OH)_3$, is a colourless, syrupy liquid with a sweet taste, insoluble in ether.

When glycerol is heated with potassium hydrogen

sulphate, or with phosphoric anhydride, acrolëin is produced and may be identified by its pungent disagreeable odour.

When a borax bead is moistened with glycerol and held near the outer edge of a Bunsen flame it gives a green colouration.

The following are crystalline solids :—

Erythritol, $C_4H_6(OH)_4$, M.P. 112° .

Oxidation with conc. nitric acid converts erythritol into oxalic acid. The tetra-benzoate melts at $186-187^\circ$.

Arabitol, $CH_2OH(CHOH)_3CH_2OH$, M.P. 102° .

Mannitol, $CH_2OH(CHOH)_4CH_2OH$, M.P. 166° .

The hexabenzoate melts at $124-125^\circ$, the hexacetate at 119° .

VI. ETHERS.

THESE are neutral, volatile liquids, practically insoluble in water and chemically very indifferent. They can be best identified by their boiling points.

Ethyl Ether, $(C_2H_5)_2O$, B.P. $34-35^\circ$.

Normal Propyl Ether, $(C_3H_7)_2O$, B.P. $90-91^\circ$.

Anisol, $C_6H_5 \cdot O \cdot CH_3$ is an ethereal smelling liquid boiling at $154-155^\circ$. Heated with concentrated hydriodic acid to 140° it decomposes into phenol and methyl iodide.

Phenetol, $C_6H_5 \cdot O \cdot C_2H_5$, boils at 172° .

Phenyl Ether, $C_6H_5 \cdot O \cdot C_6H_5$, crystallizes in long needles and possesses a smell resembling geraniums. It melts at 28° , boils at 252° . It is not reduced on heating with hydriodic acid,

VII. PHENOLS.

THE phenols, with the exception of *m*-cresol are crystalline solids. They are of an acid nature, the hydrogen of the hydroxyl being readily substituted by metals. The monohydric phenols are but sparingly soluble in water, the polyhydric phenols readily soluble.

Phenol, C_6H_5OH , is a colourless crystalline solid, which gradually acquires a reddish colour, and deliquesces on exposure to air. It melts at 43° , boils at 181° ; it dissolves in 15 parts water and gives a violet colouration in neutral solution with ferric chloride.

When an aqueous solution of phenol is mixed with one-fourth of its volume of ammonia solution and then with a few drops of a bleaching powder solution, and gently warmed, a fine blue colour is produced, which soon disappears.

Bromine water precipitates tribromophenol from solutions as an oil which soon crystallizes and melts at 92° . The crude tribromophenol may if necessary be purified by recrystallizing from alcohol. The benzoate melts at 69° .

o-Cresol, $\text{C}_6\text{H}_4 \begin{smallmatrix} \text{CH}_3 \\ \text{OH} \end{smallmatrix}$ M.P. 31° . On fusion

with potassium hydroxide, salicylic acid is formed. With picric acid, a picrate, $2\text{C}_7\text{H}_5\text{O}, 3\text{C}_6\text{H}_3\text{N}_3\text{O}_7$, melting at 88° is obtained in orange yellow needles.

m-Cresol is liquid and boils at $202\text{--}203^\circ$. Ferric chloride colours an aqueous solution of *m*-cresol deep blue violet. The benzoate melts at 54° .

p-Cresol melts at 36° and gives with ferric chloride a blue colouration. The benzoate melts at $71\text{--}72^\circ$.

Carvaerol, $\begin{smallmatrix} \text{CH}_3 \\ \text{C}_6\text{H}_7 \end{smallmatrix} \text{OH}$ is a thick oil boiling at $236\text{--}237^\circ$.

An alcoholic solution of carvaerol is coloured green on addition of ferric chloride.

Thymol, $\begin{smallmatrix} \text{CH}_3 \\ \text{C}_6\text{H}_7 \end{smallmatrix} \text{OH}$ melts at $50\text{--}51^\circ$, and possesses

a pleasant aromatic odour. An aqueous solution of thymol treated with one half its volume of glacial acetic acid and one volume of concentrated sulphuric acid produces on warming a red violet colouration.

α -Naphthol, $\text{C}_{10}\text{H}_7\text{OH}$, melts at 94° , is practically insoluble in cold water, readily soluble in alcohol and ether. Bleaching powder solution produces a dark violet colouration. A naphthol picrate is

produced, by mixing alcoholic solutions of the components, in orange needles melting at 189° .

α -Naphthol acetate melts at 46° .

PREPARATION OF ACETATE.—One gram of the naphthol is boiled for about fifteen minutes with 2 grams acetic anhydride. On pouring the product into water the crude acetate separates out. It is purified by recrystallization from a small quantity of water.

α -Naphthol benzoate melts at 56° .

β -Naphthol melts at 122° . With ferric chloride a greenish colouration is produced in an aqueous solution of the naphthol, from which after some time a white flocculent precipitate is thrown down. β -Naphthol dissolved in strong potassium hydroxide solution and treated with chloroform produces on warming to 50° a blue colouration which gradually becomes green and finally brown. β -Naphthol acetate, $C_{10}H_7OOC \cdot CH_3$, melts at 70° . The benzoate melts at 107° .

Catechol (pyrocatechin), $C_6H_4 \begin{smallmatrix} \text{OH}(1) \\ \text{OH}(2) \end{smallmatrix}$, melts at 104° . It reduces Fehling's solution on warming. With lead acetate a white precipitate is obtained. Ferric chloride produces in an aqueous solution of catechol an emerald-green colouration, which on addition of sodium bicarbonate becomes violet-red.

Gualacol, $C_6H_4 \begin{smallmatrix} \text{OCH}_3(1) \\ \text{OH}(2) \end{smallmatrix}$, melts at 31° . Ferric chloride gives its alcoholic solution an emerald-green colour.

Resorcinol, $C_6H_4 \begin{smallmatrix} \text{OH (1)} \\ \text{OH (3)} \end{smallmatrix}$, M.P. 118°. Ferric

chloride produces a dark violet colouration. If equal weights of resorcinol and phthalic anhydride are gently heated together in a test-tube a yellow melt is obtained, which, when dissolved in dilute caustic soda solution, produces a solution showing a fine green fluorescence (formation of fluorescein). Bromine water precipitates tribromoresorcinol, M.P. 111°, from an aqueous solution of resorcinol. The dibenzoate melts at 117°.

Quinol (hydroquinone), $C_6H_4 \begin{smallmatrix} \text{OH (1)} \\ \text{OH (4)} \end{smallmatrix}$, melts at 169° and is readily soluble in water, alcohol and ether.

With ferric chloride quinhydrone is readily obtained in beautiful metallic green crystals by adding ferric chloride to a concentrated solution of hydroquinone in water. The solution becomes at first red and then rapidly darkens.

The diacetate and the dibenzoate melt respectively at 123° and 199°.

Oreinol, $C_6H_3 \begin{smallmatrix} \text{CH}_3 \text{ (1)} \\ \text{OH (3)} \\ \text{OH (5)} \end{smallmatrix}$, M.P. 107°, is easily soluble

in water, alcohol and ether. On exposure to the air it becomes red. With ferric chloride a violet-black colouration is produced. Bleaching powder solution produces a dark red colouration which soon changes to yellow. The diacetate melts at 25°, the dibenzoate at 88°.

Pyrogallol, $C_6H_3(OH)_3$ 1 : 2 : 3, melts at 132–133°

and is readily soluble in water, alcohol and ether. An alkaline solution of pyrogallol quickly absorbs oxygen from the air and becomes brown. An aqueous solution of pyrogallol is turned brown on addition of nitrous acid solution. The tribenzoate melts at 90° .

Phloroglucinol, $C_6H_3(OH)_3$ 1 : 3 : 5, melts at 218° on quickly heating. It is readily soluble in water, alcohol and ether. With ferric chloride a blue-violet colouration is obtained. The acetate melts at 105° , the benzoate at 173° . The trioxime melts at 43° .

ALCOHOL PHENOLS.

Saligenin, $C_6H_4 \begin{smallmatrix} \text{OH} \\ \text{CH}_2\text{OH} \end{smallmatrix} \begin{smallmatrix} (1) \\ (2) \end{smallmatrix}$, M.P. 86° , is very readily soluble in boiling water, alcohol and ether. It dissolves in concentrated sulphuric acid with an intense red colour. With ferric chloride it gives a blue colouration.

Anisyl Alcohol, $C_6H_4 \begin{smallmatrix} \text{OCH}_3 \\ \text{CH}_2\text{OH} \end{smallmatrix} \begin{smallmatrix} (1) \\ (4) \end{smallmatrix}$, melts at 45° .

Dilute nitric acid oxidizes it to anisic acid, M.P. 185° .

HALOGEN PHENOLS.

o-Chlorophenol, $C_6H_4 \begin{smallmatrix} \text{OH} \\ \text{Cl} \end{smallmatrix}$, has an unpleasant odour and boils at $175-176^{\circ}$.

m-Chlorophenol is obtained as crystals melting at 28.5° .

p-Chlorophenol melts at 37°. It possesses a faint, unpleasant odour. Fused with alkalis it gives resorcinol.

o-Bromophenol is an unpleasant-smelling oil, boiling at 194–195°.


m-Bromophenol melts at 32–33°.

p-Bromophenol melts at 63–64°.

o-Iodophenol is obtained as needles, melting at 43°. It is fairly soluble in hot water, readily soluble in alcohol and ether. By warming with nitric acid (Sp. Gr. = 1.2) picric acid is formed.

m-Iodophenol melts at 40°.

p-Iodophenol forms long needles melting at 92°. By boiling with nitric acid it is readily converted into picric acid.



has an acid reaction. With hydrochloric acid and potassium chlorate, chloranil (*see* page 59) is formed.

Tribromophenol (2 : 4 : 6) melts at 95°. The acetate recrystallized from alcohol melts at 82°.

AMINOPHENOLS

o-Aminophenol, $C_6H_4 \begin{smallmatrix} \text{NH}_2 \\ \text{OH} \end{smallmatrix}$, when pure is colourless, but readily becomes brown. It melts at 170°, and is somewhat sparingly soluble in water, readily in ether. The acetate melts at 150°.

p-Aminophenol melts with decomposition at 184°. With chromic acid it is readily oxidized to quinone.

p-Aminophenol Methyl Ester (*pp*— anisidine)

$\text{C}_6\text{H}_4 \begin{matrix} \text{OCH}_3 & (1) \\ \text{NH}_2 & (4) \end{matrix}$ forms plates melting at 52–56°.

Phenacetin, $\text{C}_6\text{H}_4 \begin{matrix} \text{OC}_2\text{H}_5 & (1) \\ \text{NHCOCH}_3 & (4) \end{matrix}$ melts at 135°.

On hydrolysis with caustic soda it yields sodium acetate, and phenetidine, B.P. 244°.

VIII. ALDEHYDES.

THE lower aldehydes are volatile liquids soluble in water and possessing a characteristic odour. The higher members of the series are solids, insoluble in water.

In chemical properties the aldehydes are neutral substances, easily oxidized to corresponding acids, their ready oxidation by the salts of the noble metals being one of the characteristics of the aldehydes.

All aldehydes form oximes with hydroxylamine, phenylhydrazones with phenylhydrazine and semicarbazones with semicarbazide hydrochloride. As these derivatives can readily be obtained in the pure state and are generally solids, their preparation and melting-point determination serve as an aid in the identification of an aldehyde.

PREPARATION OF AN OXIME.—One gram of the

aldehyde dissolved in the least possible quantity of alcohol is treated with a slight excess of hydroxylamine hydrochloride dissolved in a little water. To this mixture is added the theoretical quantity of sodium hydroxide in concentrated solution. If any precipitate is now formed, alcohol or water, as may be necessary, is added to bring about complete solution. The solution is then heated on a water-bath under a reflux condenser for from one to two hours. The reaction mixture is then poured into water, made faintly acid with acetic acid and the oxime extracted with ether, the ethereal solution dried, the ether distilled off and the oxime recrystallized from some suitable solvent.

PREPARATION OF A PHENYLHYDRAZONE.—One cc. phenylhydrazine is added to about 5 cc. water and sufficient acetic acid added, with vigorous shaking, to bring about complete solution. About 0.5 gram of the aldehyde is now added (in solution if necessary) and the mixture vigorously shaken and very gently warmed. The phenylhydrazone separates out and, if solid, is filtered off, washed with dilute acetic acid, dried on a porous plate or between filter paper, and finally recrystallized from alcohol or benzene.

PREPARATION OF A SEMICARBAZONE.—One gram of the aldehyde, dissolved in the smallest possible quantity of alcohol is added to a solution containing 2 grams semicarbazide hydrochloride dissolved in the minimum quantity of water. A cold alcoholic solution containing 2 grams potassium acetate, best

prepared by boiling the acetate with alcohol, is now added, and, if any precipitate forms, water or alcohol, as the case requires, is then carefully added until a clear solution is obtained. The mixture is then allowed to stand for some hours, the aldehyde semicarbazone gradually separating out.

The addition of a few drops of methyl alcohol (free from acetone) often accelerates the separation of the semicarbazone. The semicarbazone obtained is finally recrystallized from methyl or ethyl alcohol.

Formaldehyde, $\text{H}\cdot\text{CHO}$, is gaseous at the ordinary temperature and is usually met with in solution, the aqueous solution possessing a penetrating, suffocating odour. When its aqueous solution is mixed with ammoniacal silver nitrate solution a silver mirror is obtained.

Trioxymethylene $(\text{H}\cdot\text{CHO})_3$, is a crystalline compound which sublimes readily under 100° . When strongly heated it is decomposed into pure gaseous formaldehyde. It is readily soluble in cold sodium hydroxide solution, insoluble in alcohol and ether.

Acetaldehyde, $\text{CH}_3\cdot\text{CHO}$, is obtained as a strongly-smelling suffocating liquid, boiling at 21° and miscible in all proportions with water. Boiling with potassium hydroxide solution produces the yellow-brown aldehyde resin. Acetaldehyde combines readily with ammonia gas to give aldehyde ammonia, a colourless crystalline solid. Its phenylhydrazone melts at 99° , the semicarbazone at 162° , the oxime at 47° .

Para-aldehyde $(\text{C}_2\text{H}_4\text{O})_3$, boils at 124° and on dis-

tillation is completely converted into acetaldehyde. It gives none of the usual aldehyde reactions.

Meta-aldehyde (C_2H_4O)₃, is isomeric with para-aldehyde. It is a crystalline solid which sublimes at 112° without melting. On distillation with dilute sulphuric acid it breaks down into acetaldehyde. Like para-aldehyde it gives none of the aldehyde reactions.

Aldol, $CH_3 \cdot CH(OH)CH_2 \cdot CHO$, is a syrupy liquid which decomposes at 135° into crotonaldehyde and water—

$CH_3CH(OH)CH_2 \cdot CHO = CH_3CH : CH \cdot CHO + H_2O$. It is soluble in water and alcohol, and gives the usual aldehyde reactions.

Acrolëin, $CH_2 : CH \cdot CHO$, is a pungent-smelling liquid, boiling at 52–53°. It is readily oxidized to acrylic acid.

Crotonaldehyde, $CH_3 \cdot CH : CH \cdot CHO$, is an extremely disagreeable-smelling liquid boiling at 104°. The oxime melts at 119–120°.


Benzaldehyde, $C_6H_5 \cdot CHO$, is a pleasant aromatic smelling liquid, which boils at 179°. It does not reduce Fehling's solution but gives most of the aldehyde reactions. Its oxime melts at 35°, the phenylhydrazone at 155°, the semicarbazone at 214°.

o-Toluic Aldehyde, $C_6H_4 \begin{matrix} \diagup CH_3 \\ \diagdown CHO \end{matrix}$, boils at 200°. Its oxime melts at 48–49°.

m-Toluic Aldehyde boils at 199°.

p-Toluic Aldehyde is a liquid with a pepper-like odour, boiling at 204°. The oxime exists in two

forms. The anti-derivative melts at 79° , the syn-compound at 109° .


p-Cumic Aldehyde, Cuminol,  is a pleasant,
 $\text{CH}(\text{CH}_3)_2$

aromatic-smelling liquid, boiling at 237° . Its oxime melts at 58° .

Cinnamic Aldehyde, $\text{C}_6\text{H}_5\cdot\text{CH}:\text{CH}\cdot\text{CHO}$, decomposes on boiling. The phenylhydrazone melts at 168° .

SUBSTITUTED ALDEHYDES.

Chloral, $\text{CCl}_3\cdot\text{CHO}$, is a colourless, pungent-smelling liquid boiling at $97\text{--}98^{\circ}$. Concentrated alkali solutions decompose chloral even at ordinary temperatures into chloroform and alkali formate—



The oxime melts at $39\text{--}40^{\circ}$.

Chloral Hydrate, $\text{CCl}_3\cdot\text{CH}(\text{OH})_2$, melts at 57° . It is readily soluble in water and alcohol. It does not show the usual aldehyde reactions. By shaking with concentrated sulphuric acid it is immediately converted into chloral.

Bromal, $\text{CBr}_3\cdot\text{CHO}$, boils at 174° . With alkalis it is decomposed into bromoform and alkali formate.

Bromal Hydrate melts at $53\text{--}54^{\circ}$.

***o*-Chlorobenzaldehyde**, $\text{C}_6\text{H}_4\text{--}\begin{matrix} \text{Cl} \\ \diagup \\ \text{CHO} \end{matrix}$, is a liquid, boiling at $213\text{--}214^{\circ}$, with a strong odour. Its oxime melts at 75° (anti), the syn. derivative at $98\text{--}102^{\circ}$.

m-Chlorobenzaldehyde is obtained in long prisms, melting at 17–18°, boiling at 213°. Oxime, M.P. 70° (syn.).

p-Chlorobenzaldehyde is a solid with an odour like benzaldehyde, melting at 47–48°.

o-Nitrobenzaldehyde, $\text{C}_6\text{H}_4 \begin{smallmatrix} \text{CHO} \\ \text{NO}_2 \end{smallmatrix}$, is obtained in the form of long, pale yellow needles, melting at 44°. It has an odour similar to that of benzaldehyde, is readily soluble in alcohol and ether, sparingly in water. Concentrated aqueous sodium hydroxide converts it readily into *o*-nitrobenzoic acid and *o*-nitrobenzylalcohol. The oxime melts at 96°.

m-Nitrobenzaldehyde melts at 58°. It is fairly soluble in hot water, readily in alcohol and ether. The oxime melts at 118°.

p-Nitrobenzaldehyde melts at 106°. With chromic acid it is converted into its corresponding acid. The oxime melts at 129°.

o-Nitrocinnamic aldehyde, $\text{C}_6\text{H}_4 \begin{smallmatrix} \text{CH:CH:CHO} \\ \text{NO}_2 \end{smallmatrix}$, melts at 127° and is soluble in hot water.

m-Nitrocinnamic aldehyde, M.P. 116°.

p-Nitrocinnamic aldehyde melts at 141°. The oxime melts at 178–179°.

o-Aminobenzaldehyde, $\text{C}_6\text{H}_4 \begin{smallmatrix} \text{NH}_2 \\ \text{CHO} \end{smallmatrix}$, is obtained in silvery leaflets melting at 39–40°, readily soluble in alcohol and ether, sparingly in water. The acetyl derivative melts at 70°, the oxime at 132°.

***m*-Aminobenzaldehyde** is a yellow amorphous substance. The oxime melts at 88°.

***p*-Aminobenzaldehyde** melts at 70–71°. The hydrochloride is obtained in red crystals. The acetyl derivative melts at 155°, the oxime at 124°.

Salicylic Aldehyde, $C_6H_4 \begin{matrix} \text{OH} \\ \text{CHO} \end{matrix} \begin{matrix} (1) \\ (2) \end{matrix}$, is a pleasant-smelling oil, boiling at 196°. On oxidation it yields salicylic acid. The acetyl derivative melts at 37°, the oxime at 57°.

***m*-Hydroxybenzaldehyde** melts at 104°. The aqueous solution is coloured deep violet with ferric chloride solution. The oxime melts at 87°.


***p*-Hydroxybenzaldehyde** melts at 115–116° and sublimes undecomposed. The aqueous solution is coloured faintly violet with ferric chloride. The oxime melts at 65°.

Anisic Aldehyde, $C_6H_4 \begin{matrix} \text{CHO} \\ \text{OCH}_3 \end{matrix} \begin{matrix} (1) \\ (4) \end{matrix}$, boils at 248°. The oxime melts at 61°.

Protocatechuic Aldehyde, $C_6H_3 \begin{matrix} \text{CHO} \\ \text{OH} \\ \text{OH} \end{matrix} \begin{matrix} (1) \\ (3) \\ (4) \end{matrix}$, melts at 153°. Ferric chloride colours an aqueous solution of protocatechuic aldehyde green; on addition of sodium carbonate the colour changes to violet and then to red. The oxime melts at 150°.

Vanillin, $C_6H_3 \begin{matrix} \text{CHO} \\ \text{OCH}_3 \\ \text{OH} \end{matrix} \begin{matrix} (1) \\ (3) \\ (4) \end{matrix}$ has a pleasant, vanilla-like odour and melts at 80–81°. It gives with ferric

chloride a blue colouration. The benzoate melts at 75° , the oxime at 117° .

Piperonal, C_6H_5 , has a pleasant helio-

trope odour. It melts at 37° . The oxime melts at $110-112^{\circ}$, the phenylhydrazone at 100° .

Closely related to the aldehydes themselves are the aldehyde ethers. These are liquids, sparingly soluble in water, which on boiling with dilute hydrochloric acid are readily broken up into their constituent aldehyde and alcohol.

Methylal $\text{CH}_2(\text{OCH}_3)_2$, B.P. 42° .

Acetal, $\text{CH}_3\cdot\text{CH}(\text{OC}_2\text{H}_5)_2$, B.P. 104° .

IX. KETONES.

THE aliphatic ketones are liquids, the common aromatic ketones are solids. They are only oxidized with difficulty and therefore do not reduce alkaline silver solutions. Only those ketones with a methyl group directly attached to the carbonyl group form bisulphite compounds. Like aldehydes they form oximes, phenylhydrazones, and semicarbazones. The methods for the preparation of these derivatives are similar to those employed in the case of the aldehydes.

Acetone, $\text{CH}_3\cdot\text{CO}\cdot\text{CH}_3$, B.P. 56° , is miscible with

water in all proportions. With iodine and caustic potash it forms iodoform. The oxime melts at 59–60°.

Methylethylketone, $\text{CH}_3 \cdot \text{CO} \cdot \text{C}_2\text{H}_5$, boils at 80–81°. The oxime is an oil, the semicarbazone melts at 135–136°.

Diethylketone, $\text{C}_2\text{H}_5 \cdot \text{CO} \cdot \text{C}_2\text{H}_5$, B.P. 102–103°.

Dipropylketone, $\text{C}_3\text{H}_7 \cdot \text{CO} \cdot \text{C}_3\text{H}_7$, B.P. 144°.

Pinacolone, $\begin{array}{c} \text{CH}_3 \\ \diagup \\ \text{CH}_3 \end{array} \text{C} \cdot \text{CO} \cdot \text{CH}_3$, is a peppermint-smelling liquid boiling at 106°. It forms no bisulphite compound. The oxime melts at 74–75°.

Acetophenone, $\text{C}_6\text{H}_5 \cdot \text{CO} \cdot \text{CH}_3$, melts at 20°. With picric acid it gives a picrate, greenish-yellow crystals melting at 53°. The oxime melts at 59°.

Benzophenone, $\text{C}_6\text{H}_5 \cdot \text{CO} \cdot \text{C}_6\text{H}_5$, melts at 48–49°. The oxime melts at 139–140°, the phenyl-hydrazone at 137°.

Benzoin, $\text{C}_6\text{H}_5 \cdot \text{CHOH} \cdot \text{CO} \cdot \text{C}_6\text{H}_5$, melts at 137°. It is converted into benzil by means of nitric acid. Fehling's solution is readily reduced by means of benzoin. The oxime melts at 151–152° (*a*-derivative).

Deoxybenzoin, $\text{C}_6\text{H}_5 \cdot \text{CH}_2 \cdot \text{CO} \cdot \text{C}_6\text{H}_5$, melts at 60°. The oxime melts at 98°, the phenylhydrazone at 116°.

Benzil, $\text{C}_6\text{H}_5 \cdot \text{CO} \cdot \text{CO} \cdot \text{C}_6\text{H}_5$, melts at 95°. The dioxime melts with decomposition at 237°.

X. ACIDS.

To obtain the acid from a salt, the latter is dissolved in water (if it is insoluble, it is boiled with sodium carbonate and filtered), and the solution acidified. In many cases the acid will be precipitated; if soluble in water it is extracted with ether, the ethereal solution dried and the ether distilled off. The physical and chemical properties of the acid are then determined.

1. SATURATED ALIPHATIC MONOBASIC ACIDS.

The lower members are liquids, soluble in water, the solubility decreasing with increase of molecular weight. The higher members are odourless solids, insoluble in water.

Formic Acid, $\text{H}\cdot\text{COOH}$, when anhydrous boils at 101° . By the action of concentrated sulphuric acid, carbon monoxide is evolved. Formic acid reduces solutions of silver and mercury salts.

Acetic Acid, $\text{CH}_3\cdot\text{COOH}$, when free from water melts at 16.5° , and boils at 118° . The addition of ferric chloride to a neutral solution of an acetate produces a deep red colour, which on warming disappears, while a brown flocculent precipitate of basic ferric acetate is thrown down.

When acetic acid is warmed with alcohol and concentrated sulphuric acid, ethyl acetate is formed, which may be recognized by its pleasant fruity odour.

Propionic Acid, $\text{C}_2\text{H}_5\cdot\text{COOH}$, boils at 141° . It is

precipitated as an oil from aqueous solution by the addition of calcium chloride. The ethyl ester boils at 98-99°.

n-Butyric Acid, $\text{CH}_3\cdot\text{CH}_2\cdot\text{CH}_2\cdot\text{COOH}$, boiling at 162°, has an extremely unpleasant odour. Its ethyl ester boils at 120°.

Isobutyric Acid, $\begin{array}{c} \text{CH}_3 \\ \text{CH}_3 \end{array} > \text{CH}\cdot\text{COOH}$, boils at 155° and is insoluble in water. The ethyl ester boils at 110°.

Palmitic Acid, $\text{CH}_3\cdot(\text{CH}_2)_{14}\cdot\text{COOH}$, melts at 62°. The lead salt formed by adding lead acetate to a neutral solution of ammonium palmitate melts at 112°.

Stearic Acid, $\text{CH}_3\cdot(\text{CH}_2)_{16}\cdot\text{COOH}$, melts at 69.2° and the lead salt at 125°.

2. SATURATED ALIPHATIC POLYBASIC ACIDS.

These acids are solids, and are soluble in water.

Oxalic Acid, $\begin{array}{c} \text{COOH} \\ | \\ \text{COOH} \end{array} + 2\text{H}_2\text{O}$, melts at 98°. When

acted on by concentrated sulphuric acid, carbon monoxide and carbon dioxide are evolved. Potassium permanganate reacts with oxalic acid, with formation of carbon dioxide and water. The calcium salt is insoluble in water and in acetic acid. The dimethyl ester, which can be prepared by boiling the anhydrous acid with methyl alcohol for about ten minutes, is a solid melting at 54°.

Malonic Acid, $\text{CH}_2 \begin{matrix} \diagup \text{COOH} \\ \diagdown \text{COOH} \end{matrix}$, melts at 132° . On heating more strongly, it is decomposed into acetic acid and carbon dioxide.

Succinic Acid, $\begin{matrix} \text{CH}_2 \cdot \text{COOH} \\ | \\ \text{CH}_2 \cdot \text{COOH} \end{matrix}$, melts at 185° . When

a neutral solution of ferric chloride is added to a neutral solution of a succinate, a brown-red precipitate of basic ferric succinate is produced. Succinic

anhydride $\begin{matrix} \text{CH}_2\text{CO} \diagdown \\ | \\ \text{CH}_2\text{CO} \diagup \end{matrix} \text{O}$ melting at 119° , is obtained

by the action of acetyl chloride on the acid at 50° . It is purified by recrystallizing from chloroform.


Glutaric Acid, $\text{CH}_2 \begin{matrix} \diagup \text{CH}_2\text{COOH} \\ \diagdown \text{CH}_2\text{COOH} \end{matrix}$, melting at 97.5° , is converted into the anhydride (M.P. $56-57^\circ$) by heating for some time.

3. UNSATURATED ALIPHATIC ACIDS.


Acrylic Acid, $\text{CH}_2\text{:CH}\cdot\text{COOH}$, boils at 140° and polymerizes. The lead salt crystallizes in fine needles. When acrylic acid is fused with potassium hydroxide, it is decomposed into potassium formate, potassium acetate and hydrogen. The ethyl ester boils at $101-102^\circ$.

Crotonic Acid, $\text{CH}_3\text{:CH:CH}\cdot\text{COOH}$, melting at $71-72^\circ$, when fused with potassium hydroxide produces potassium acetate and hydrogen. Its ethyl ester boils at 142° .

Oleic Acid, $\text{CH}_3 \cdot (\text{CH}_2)_7 \cdot \text{CH} : \text{CH} \cdot (\text{CH}_2)_7 \cdot \text{COOH}$, melts at 14° , and boils at 223° at 10 mm. The lead salt is soluble in ether and melts about 80° . When oleic acid is shaken with nitrous acid, it is transformed into the isomeric **Elaidic Acid**, melting at $44\text{--}45^\circ$.



200° , without melting. When distilled with phosphorus pentoxide, maleic anhydride (M.P. 60°) is formed. By the action of bromine at 100° on fumaric acid, dibromsuccinic acid, melting at $255\text{--}256^\circ$ with decomposition, is produced.



converted into the anhydride (M.P. 60°) by heating under diminished pressure at 100° .

4. AROMATIC ACIDS.

Benzoic Acid, $\text{C}_6\text{H}_5 \cdot \text{COOH}$, melts at 121° , and can be sublimed easily. It is practically insoluble in cold water, easily soluble in alcohol and ether. When heated with lime, benzene is produced. Ethyl benzoate (B.P. 213°), a pleasant-smelling liquid, is easily formed by heating together benzoic acid, ethyl alcohol and a small quantity of concentrated sulphuric acid.



soluble in hot water. Dilute nitric acid oxidizes it to phthalic acid.



m-Toluic Acid, M.P. 110° , is readily soluble in water.

p-Toluic Acid, M.P. $176-177^{\circ}$. The methyl ester, a strongly smelling solid, melts at 32° .

α -Naphthole Acid, $C_{10}H_7 \cdot COOH$, M.P. 160° and

β -Naphthole Acid, M.P. 182° are converted into naphthalene by distillation with lime. When the α -compound is oxidized with chromic acid in glacial acetic acid, phthalic acid is produced. Methyl β -naphthoate is a solid melting at 77° .

Phthalic Acid, $C_6H_4 \begin{matrix} \swarrow COOH (1) \\ \searrow COOH (2) \end{matrix}$, on heating forms phthalic anhydride (M.P. 128°). When the latter is heated with phenol and a little anhydrous zinc chloride, phenolphthalein is produced—



Phenylacetic Acid, $C_6H_5 \cdot CH_2 \cdot COOH$, M.P. 76° , is sparingly soluble in cold water.

Hydrocinnaminic Acid, $C_6H_5 \cdot CH_2 \cdot CH_2 \cdot COOH$, M.P. 48° .

Cinnamic Acid, $C_6H_5 \cdot CH : CH \cdot COOH$, M.P. 133° . The methyl ester melts at 33° .

5. SUBSTITUTED ACIDS.

A. HYDROXY ACIDS.

Glycollic Acid, $\text{CH}_2\text{OH} \cdot \text{COOH}$, melting at $78-79^\circ$, when heated to 100° is converted into an anhydride, M.P. $128-130^\circ$.

Lactic Acid, $\text{CH}_3 \cdot \text{CHOH} \cdot \text{COOH}$, is a syrup, easily soluble in water and alcohol, difficultly soluble in ether. Hydriodic Acid reduces it to propionic acid, while at the same time free iodine is produced. The silver salt melts at 100° .

Malic Acid, $\begin{array}{c} \text{CHOH} \cdot \text{COOH} \\ | \\ \text{CH}_2 \cdot \text{COOH} \end{array}$, melts at 100° . When calcium chloride and alcohol are added to a neutral solution of a malate, calcium malate is precipitated.

Tartaric Acid, $\begin{array}{c} \text{CHOH} \cdot \text{COOH} \\ | \\ \text{CHOH} \cdot \text{COOH} \end{array}$, melting at $168-170^\circ$, is decomposed on heating strongly, with production of a smell of burnt sugar. Calcium chloride added to a neutral solution of a tartrate (but not of tartaric acid) precipitates calcium tartrate. When ammoniacal silver nitrate solution is added to a neutral solution of a tartrate, a silver mirror is produced on warming.

Citric Acid, $\begin{array}{c} \text{CH}_2 \cdot \text{COOH} \\ | \\ \text{C}(\text{OH}) \cdot \text{COOH} \\ | \\ \text{CH}_2 \cdot \text{COOH} \end{array} + \text{H}_2\text{O}$, is easily soluble in water and alcohol. It loses its water of crystalliz-

ation at 130° , and melts at 153° . When calcium chloride is added to a neutral solution of a citrate and the mixture heated for some time, calcium citrate is precipitated.

Salicylic Acid, $C_6H_4 \begin{matrix} \swarrow OH \\ \searrow COOH \end{matrix} \begin{matrix} (1) \\ (2) \end{matrix}$, melting at 155° , is converted into phenol when it is heated with lime. Ferric chloride added to a solution of salicylic acid produces a deep violet colouration, which is not destroyed by addition of acetic acid. Methyl salicylate, which has a characteristic odour, boils at 224° .

Protocatecholic Acid, $C_6H_3 \begin{matrix} \swarrow OH \\ \searrow OH \\ \searrow COOH \end{matrix} \begin{matrix} (1) \\ (2) \\ (4) \end{matrix}$, melts at 199° .

Ferric chloride added to a solution of the acid, produces a green colouration, which, on addition of sodium hydroxide, changes to blue and finally to red.

Gallie Acid, $C_6H_2(OH)_3 \cdot COOH + H_2O$, decomposes at 220° , into carbon dioxide and pyrogallol (see page 26). Ferric chloride produces a blue-black precipitate, soluble in hydrochloric acid but reprecipitated by ammonia.

Tannic Acid is an amorphous powder, more soluble in water than in alcohol. Ferric chloride solution added to a solution of the acid, produces a blue-black colouration. When sulphuric acid is added to a concentrated solution of tannic acid, a white precipitate is formed, which in contact with air turns blue.

Mandelic Acid, $C_6H_5 \cdot CHOH \cdot COOH$, melts at 118° ,

and is easily soluble in water. Oxidation with nitric acid produces benzaldehyde. The methyl ester melts at 52°.

B. HALOGEN SUBSTITUTED ACIDS.

Monochloroacetic Acid, $\text{CH}_2\text{Cl}\cdot\text{COOH}$, melting at 62–63°, heated with caustic potash forms glycollic acid.

Dichloroacetic Acid, $\text{CHCl}_2\cdot\text{COOH}$, is a liquid, boiling at 189–191°.

Trichloroacetic Acid, $\text{CCl}_3\cdot\text{COOH}$, melts at 55°. When digested with caustic potash, chloroform and potassium carbonate are formed.

Monobromoacetic Acid, $\text{CH}_2\text{Br}\cdot\text{COOH}$, melts at 49–50°.

Dibromoacetic Acid, $\text{CHBr}_2\cdot\text{COOH}$ melts at 48°.

Tribromoacetic Acid, $\text{CBr}_3\cdot\text{COOH}$ melts at 135°.

o-Chlorobenzoic Acid, $\text{C}_6\text{H}_4\begin{smallmatrix} \text{Cl} \\ \diagup \\ \text{COOH} \end{smallmatrix}$, M.P. 137°.

Reduction with sodium amalgam produces benzoic acid.

m-Chlorobenzoic Acid, M.P. 153°.

p-Chlorobenzoic Acid, M.P. 236°.

o-Bromobenzoic Acid, $\text{C}_6\text{H}_4\begin{smallmatrix} \text{Br} \\ \diagup \\ \text{COOH} \end{smallmatrix}$, M.P. 147°.

m-Bromobenzoic Acid, M.P. 155°.

p-Bromobenzoic Acid, M.P. 251°.

When the halogen benzoic acids are distilled with lime, the corresponding halogen derivatives of benzene are produced.

C. NITRO ACIDS.

o-Nitrobenzoic Acid, $C_6H_4 \begin{smallmatrix} \text{NO}_2 \\ \text{COOH} \end{smallmatrix}$, M.P. 147°.

m-Nitrobenzoic Acid, M.P. 141°.

p-Nitrobenzoic Acid, M.P. 238°.

These acids can be reduced to the corresponding amino-acids by tin and hydrochloric acid.

o-Nitrocinnamic Acid, $C_6H_4 \begin{smallmatrix} \text{NO}_2 \\ \text{CH} : \text{CH} \cdot \text{COOH} \end{smallmatrix}$,
M.P. 237-240°.

m-Nitrocinnamic Acid, light yellow needles. M.P. 197°.

p-Nitrocinnamic Acid, M.P. 286°.

Oxidation with alkaline potassium permanganate solution converts these acids into the corresponding nitrobenzaldehyde and nitrobenzoic acid.

D. AMINO ACIDS.

Aminoacetic Acid (glycine), $CH_2NH_2 \cdot COOH$, M.P. 232°, with decomposition, is converted by nitrous acid into glycollic acid. Ferric chloride added to a solution of glycine causes an intense red colouration which is discharged by acids, and restored by ammonia.

Methyl Glycine (sarcosine), $CH_2NH(CH_3) \cdot COOH$, M.P. 210-215°, gives by the action of nitrous acid a nitroso-compound. When heated with soda-lime, sarcosine produces methylamine.

Aceturic Acid (Acetoglycine),




melts at 206°. On boiling with acids, it is decomposed into glycine and acetic acid.

Hippuric Acid (Benzoylglycine),



melting at 187°, is decomposed by boiling with dilute acids into benzoic acid and glycine.




Aminopropionic Acid (Alanine) $\text{CH}_3 \cdot \text{CHNH}_2 \cdot \text{COOH}$.

When heated with concentrated phosphoric acid solution it is decomposed into carbon dioxide, ammonia and acetaldehyde.

Aspartic Acid, $\text{COOH} \cdot \text{CH}_2 \cdot \text{CHNH}_2 \cdot \text{COOH}$. Nitrous acid converts it into malic acid.

Asparagine, $\text{CONH}_2 \cdot \text{CH}_2 \cdot \text{CHNH}_2 \cdot \text{COOH}$, forming large rhombic crystals, decomposes over 200° without melting. When heated with alkalis, ammonia and aspartic acid are produced. With nitrous acid it gives malic acid.

***o*-Aminobenzoic Acid** (Anthranilic Acid),



heated, decomposes into carbon dioxide and aniline. Nitrous acid transforms it into salicylic acid.

Acetylanthranilic Acid, $\text{C}_6\text{H}_4 \begin{array}{l} \text{NHCOCH}_3 \\ \text{COOH} \end{array}$, M.P. 185°.

Benzoylanthranilic Acid, $\text{C}_6\text{H}_4 \begin{array}{l} \text{NHCOC}_6\text{H}_5 \\ \text{COOH} \end{array}$, M.P.

177°.

***m*-Aminobenzoic Acid** forms small reddish crystals. M.P. 173–174°.

p-Aminobenzoic Acid, M.P. 186–187°.


o-Aminocinnamic Acid, $C_6H_4 \begin{matrix} \text{NH}_2 \\ \text{CH} : \text{CH} \cdot \text{COOH} \end{matrix}$,
M.P. 158–159°, forms yellow needles and gives
fluorescent solutions in ether and alcohol.

XI. AROMATIC SULPHONIC ACIDS.

THESE are, as a rule, very soluble in water and crystallize with difficulty. To identify them, it is best to prepare the sulphonamide and take its melting point. The sodium salts are easily obtained by salting out.

PREPARATION OF THE AMIDE.—One gram of the sodium salt and 2 grams phosphorus pentachloride are heated on a water bath. After the reaction is finished, the mixture is poured into water and the oily acid chloride extracted by means of ether. One gram of the oil and 2 grams of ammonium carbonate are heated on a water bath until the smell of the chloride disappears. The reaction product is then poured into water. After filtering, the amide is recrystallized from water and its melting-point determined.

When a sulphonic acid or a salt is fused with alkali, a phenol is produced—



METHOD.—One gram of the acid or salt is mixed

with 2 grams solid caustic potash and heated in a nickel crucible to about 200° for one hour. After cooling the product is dissolved in water, the solution acidified and the phenol extracted with ether and identified.

By passing superheated steam into the sulphonic acid, a hydrocarbon results.

Benzene sulphonic acid, $C_6H_5SO_3H$, M.P. 50°. M.P. of amide, 150°.

o-Toluene sulphonic acid, $C_6H_4\begin{smallmatrix} \text{CH}_3 \\ \text{SO}_3H \end{smallmatrix} \cdot 2H_2O$, deliquescent leaflets. M.P. of amide, 155°.

m-Toluene sulphonic acid, $C_6H_4\begin{smallmatrix} \text{CH}_3 \\ \text{SO}_3H \end{smallmatrix} \cdot H_2O$, deliquescent needles. M.P. of amide, 107°.

p-Toluene Sulphonic acid, $C_6H_4\begin{smallmatrix} \text{CH}_3 \\ \text{SO}_3H \end{smallmatrix} \cdot 4H_2O$, crystallizes in leaflets. M.P. 92°. M.P. of amide, 137°.

o-Benzene disulphonic acid, $C_6H_4\begin{smallmatrix} \text{SO}_3H \\ \text{SO}_3H \end{smallmatrix}$, M.P. of chloride, 105°; of amide, 233°.

m-Benzene disulphonic acid, M.P. of chloride, 53°; of amide, 228°.

p-Benzene disulphonic acid, M.P. of chloride, 131°; of amide, 288°.

o-Phenolsulphonic acid, $C_6H_4\begin{smallmatrix} OH \\ \text{SO}_3H \end{smallmatrix} \cdot \frac{3}{4}H_2O$, melts above 50°. When the potassium salt is treated with benzoyl chloride, potassium chloride is precipitated, and on extraction with ether, phenyl benzoate is obtained.

***m*-Phenolsulphonic acid**, $\text{C}_6\text{H}_4 \begin{smallmatrix} \text{OH} \\ \text{SO}_3\text{H} \end{smallmatrix} \cdot 2\text{H}_2\text{O}$, crystallizes in fine needles. Ferric chloride produces in solutions of the acid, a violet colouration.

***p*-Phenolsulphonic Acid** is oxidized by manganese dioxide and sulphuric acid to quinone. When boiled with hydriodic acid, *p*-phenolsulphonic acid is decomposed into phenol and sulphuric acid.

***o*-Chlorobenzenesulphonic acid**, $\text{C}_6\text{H}_4 \begin{smallmatrix} \text{Cl} \\ \text{SO}_3\text{H} \end{smallmatrix}$, The chloride melts at 28.5° ; the amide at 188° .

***m*-Chlorobenzenesulphonic acid**. The chloride is an oil; the amide melts at 148° .

***p*-Chlorobenzenesulphonic acid**. M.P. of chloride, 53° ; of amide, 143° .

***o*-Bromobenzenesulphonic acid**, $\text{C}_6\text{H}_4 \begin{smallmatrix} \text{Br} \\ \text{SO}_3\text{H} \end{smallmatrix}$. M.P. of chloride, 51° ; of amide, 186° .

***m*-Bromobenzenesulphonic acid**. The chloride is oily. M.P. of amide, 154° .

***p*-Bromobenzenesulphonic acid**, melts at 88° . M.P. of chloride, 75° ; of amide, 166° .

***o*-Iodobenzenesulphonic acid**, $\text{C}_6\text{H}_4 \begin{smallmatrix} \text{I} \\ \text{SO}_3\text{H} \end{smallmatrix}$. M.P. of chloride, 51° ; of amide, 170° .

***p*-Iodobenzenesulphonic acid**. The chloride melts at $86-87^\circ$; the amide at 183° .

***o*-Nitrobenzenesulphonic Acid**, $\text{C}_6\text{H}_4 \begin{smallmatrix} \text{NO}_2 \\ \text{SO}_3\text{H} \end{smallmatrix}$. M.P. of chloride, 67° ; of amide, 186° .

***m*-Nitrobenzenesulphonic acid**. M.P. of chloride, 60.5° ; of amide, 161° .

***p*-Nitrobenzenesulphonic acid.** The chloride is oily ; the amide melts at 181°.

***p*-Aminobenzenesulphonic acid** (Sulphanilic acid), $C_6H_4 \begin{smallmatrix} \text{NH}_2 \\ \text{SO}_3H \end{smallmatrix}$, when oxidized gives quinone. Fusion with caustic potash produces aniline and not aminophenol.

***o*-Sulphobenzole acid**, $C_6H_4 \begin{smallmatrix} \text{SO}_3H \\ \text{COOH} \end{smallmatrix}$, when anhydrous melts at 250°. On fusion with caustic potash it is converted into salicylic acid. The imino derivative, $C_6H_4 \begin{smallmatrix} \text{SO}_2 \\ \text{CO} \end{smallmatrix} \text{NH}$, known as **Saccharin**, melts with partial decomposition at 220°. By heating with alkalis, saccharin is decomposed into ammonia and *o*-sulphobenzole acid. The sodium salt,

$C_6H_4 \begin{smallmatrix} \text{SO}_2 \\ \text{CO} \end{smallmatrix} \text{NNa}$ (the saccharin of commerce) is readily soluble in water.

***α*-Naphthalene sulphonic acid**, $C_{10}H_7SO_3H \cdot H_2O$, M.P. 85–90°. The chloride crystallizes from ether in leaflets, melting at 66°. The amide melts at 150°. Potassium permanganate in acid solution oxidizes the sulphonic acid to phthalic acid.

***β*-Naphthalene sulphonic acid**, M.P. 161°. The chloride melts at 76° ; the amide at 212°.

XII. ACID ANHYDRIDES.

THE aliphatic acid anhydrides are mostly colourless liquids, insoluble in water, readily soluble in alcohol and ether, with boiling points higher than those of the corresponding acids. The aromatic acid anhydrides are solids with melting-points lower than those of the corresponding acids. Water slowly hydrolyses anhydrides with formation of acids. With alkalis they readily form salts of the corresponding acids, and with alcohols, esters are produced.

Acetic anhydride, $(\text{CH}_3\text{CO})_2\text{O}$, B.P. $137-138^\circ$.



Succinic Anhydride, $\begin{array}{c} \text{CH}_2\text{CO} \\ | \\ \text{CH}_2\text{CO} \end{array} \text{O}$, M.P. $119-120^\circ$,

reverts to the acid in moist air.

Benzole anhydride, $(\text{C}_6\text{H}_5\text{CO})_2\text{O}$, M.P. 42° .

Phthalic anhydride, $\text{C}_6\text{H}_4 \begin{array}{c} \text{CO} \\ \diagup \quad \diagdown \\ \text{CO} \end{array} \text{O}$, melts at 128°

and sublimes readily in long needles. When fused with resorcinol, fluorescein is produced (*see* p. 26).



XIII. ACID HALIDES.

THESE are pungent-smelling liquids, readily converted by water into the acid, or by alkali into the salt of the acid.

Acetyl chloride, $\text{CH}_3\cdot\text{COCl}$, B.P. 55° .

Acetyl bromide, B.P. 81° .

Acetyl iodide, B.P. 108° .

Propionyl chloride, $\text{C}_2\text{H}_5\cdot\text{COCl}$, B.P. 80° .

Propionyl bromide, B.P. 104° .

Propionyl iodide, B.P. $127\text{--}128^\circ$.

Malonyl chloride, $\text{CH}_2 \begin{matrix} \nearrow \text{COCl} \\ \searrow \text{COCl} \end{matrix}$, B.P. 58° .

Succinyl chloride, $\begin{matrix} \text{CH}_2\cdot\text{COCl} \\ | \\ \text{CH}_2\cdot\text{COCl} \end{matrix}$, B.P. $190\text{--}192^\circ$.

Benzoyl chloride, $\text{C}_6\text{H}_5\cdot\text{COCl}$, B.P. 198° .

Benzoyl bromide, B.P. $218\text{--}219^\circ$.

Phthalyl chloride, $\text{C}_6\text{H}_4 \begin{matrix} \nearrow \text{COCl} \\ \searrow \text{COCl} \end{matrix}$, B.P. 275° .

XIV. ACID AMIDES.

THESE are well crystallized substances with definite melting points. They are, as a general rule, less soluble in water than the ammonium salts of the corresponding acids. When heated with alkalies, they give ammonia and the salt of the acid. With

nitrous acid they yield nitrogen and the corresponding acid.

Acetamide, $\text{CH}_3\cdot\text{CONH}_2$, M.P. $82-83^\circ$.

Propionamide, $\text{C}_2\text{H}_5\cdot\text{CONH}_2$, M.P. 79° .

Acetobromamide, $\text{CH}_3\cdot\text{CONHBr} + \text{H}_2\text{O}$, M.P. $70-80^\circ$, when boiled with caustic potash gives potassium bromide, carbon dioxide and methylamine. The anhydrous compound melts at 108° .

CONH_2

Oxamide, $\begin{array}{c} | \\ \text{CONH}_2 \end{array}$, sublimes on heating. It is insoluble in water and in alcohol.

CONH_2

Oxamic acid, $\begin{array}{c} | \\ \text{COOH} \end{array}$, M.P. 210° with decomposition.

CONH_2

Oxamethane, $\begin{array}{c} | \\ \text{COOC}_2\text{H}_5 \end{array}$, M.P. $114-115^\circ$, when boiled with caustic potash yields ammonia, ethyl alcohol and potassium oxalate.

$\text{CH}_2\cdot\text{CONH}_2$

Succinamide, $\begin{array}{c} | \\ \text{CH}_2\cdot\text{CONH}_2 \end{array}$, at 200° is converted into the imide and ammonia.

Cyanamide, $\text{CN}\cdot\text{NH}_2$, M.P. 40° , with copper sulphate gives a black compound, copper cyanamide, CN_2Cu .¹



Benzamide, $\text{C}_6\text{H}_5\cdot\text{CONH}_2$, M.P. 128° , is soluble in hot water, alcohol and ether.

¹ Calcium cyanamide CN_2Ca is a commercial product. On addition of water, ammonia is evolved.

Phthalic Diamide, $C_6H_4 \begin{smallmatrix} \diagup CONH_2 \\ \diagdown CONH_2 \end{smallmatrix}$, on heating changes to phthalimide with loss of ammonia.

XV. ACID IMIDES.

THE imides, like the acid amides, are hydrolysed by boiling with alkalies. The only common imides are the following :—



Alcoholic caustic potash precipitates the potassium salt, $C_6H_4 \begin{smallmatrix} \diagup CO \\ \diagdown CO \end{smallmatrix} \rangle NK$, from a solution of phthalimide in alcohol. This salt is insoluble in alcohol and ether, and sparingly soluble in water.

XVI. ACID ANILIDES.

THESE are hydrolysed by heating with concentrated hydrochloric acid or alkali and the aniline distilled with steam from an alkaline solution. The acid is then identified in the non-volatile residue.

Formanilide, $H \cdot CONHC_6H_5$, melting at 46°, is

readily soluble in water, alcohol and ether. With concentrated caustic soda solution there is produced a crystalline sodium compound, $\text{H} \cdot \text{CONNaC}_6\text{H}_5$.

Acetanilide (antifebrin) $\text{CH}_3 \cdot \text{CONHC}_6\text{H}_5$, M.P. 112° .

COOH

Oxanilic acid, $\left| \begin{array}{c} \text{COOH} \\ \text{CONHC}_6\text{H}_5 \end{array} \right.$ $\cdot \text{H}_2\text{O}$, M.P. (anhydrous)

$149\text{--}150^\circ$.

CONHC_6H_5

Oxanilide, $\left| \begin{array}{c} \text{CONHC}_6\text{H}_5 \\ \text{CONHC}_6\text{H}_5 \end{array} \right.$, M.P. 245° .

Benzanilide, $\text{C}_6\text{H}_5 \cdot \text{CONHC}_6\text{H}_5$, M.P. $161\text{--}162^\circ$.

XVII. ESTERS.

THESE are mostly pleasant-smelling, volatile liquids, insoluble in water. (Methyl oxalate and methyl tartrate are solids.)

A small quantity of the ester is hydrolysed with 10 per cent. alkali solution, under a reflux condenser. Two-thirds of the liquid are distilled over, dehydrated with potassium carbonate and examined for the alcohol. The residue in the distilling flask contains the alkali salt of the acid.

Formates -methyl, B.P. 32.5° .

-ethyl, B.P. 54.5° .

-propyl, B.P. $82\text{--}83^\circ$.

-butyl, B.P. 107° .

-n-amyl, B.P. 130.5° .

Acetates -methyl, B.P. 57.5°.

-ethyl, B.P. 77°.

-propyl, B.P. 101°.

-butyl, B.P. 125°.

-isobutyl, B.P. 116°.

-*n*-amyl, B.P. 148°.

Glycol monoacetate, B.P. 182°.

Glycol diacetate, B.P. 186-187°.

Glycerol triacetate, B.P. 258-259°.

Propionates-methyl, B.P. 79.5°.

-ethyl, B.P. 99°.

Butyrates-methyl, B.P. 102.3°.

-ethyl, B.P. 120°.

Oxalates-dimethyl, M.P. 54°.

-diethyl, B.P. 186°.

Malonates-dimethyl, B.P. 181-182°.

-diethyl, B.P. 198°.

Succinates-dimethyl, M.P. 18°, B.P. 195°.

-ethyl, B.P. 216°.

Malates-dimethyl, B.P. 122° at 12 mm.

-diethyl, B.P. 129° at 12 mm.

-*n*-dipropyl, B.P. 150° at 12 mm.

Ethereal malates on heating under ordinary pressure decompose into the corresponding fumarates.

Tartrates -dimethyl, M.P. 48°.

-diethyl, B.P. 280°.

-*n*-dipropyl, B.P. 303°.

Benzoates -methyl, B.P. 199°.

-ethyl, B.P. 213°.

-*n*-propyl, B.P. 229°.

Glycol di-benzoate, M.P. 73°.

Glycerol tri-benzoate, M.P. 76°.

Salicylates -methyl, B.P. 224°.

-ethyl, B.P. 231.5°. When distilled with barium oxide ethyl phenyl ether is produced.

-phenyl (salol), M.P. 42-43°.

Phthalates -dimethyl, B.P. 282°.

-diethyl, B.P. 288°.

-diphenyl, M.P. 70°.

Cinnamates -methyl, M.P. 36°, B.P. 260°.

-ethyl, B.P. 271°.

phenyl, M.P. 72°. On distillation in air, carbon dioxide and stilbene are formed.

XVIII. QUINONES.

THESE compounds are characterized by their colour (yellow or red), peculiar odour and volatility in steam. With the exception of anthraquinone and phenanthraquinone, they are readily reduced by sulphur dioxide to the corresponding dihydric phenols.

Benzoquinone (quinone), $C_6H_4O_2$, M.P. 116°, is soluble in water. With phenol, quinone forms phenoquinone, $C_6H_4O_2 \cdot 2C_6H_5OH$, crystallizing in red needles, melting at 71°. Phenoquinone is turned

blue by caustic potash, green by barium hydroxide. Quinone liberates iodine from potassium iodide.

o-Quinone forms bright red crystals. It has no smell and is not volatile. It is reduced by sulphurous acid to catechol.

α -Naphthoquinone, $C_{10}H_6O_2$, M.P. 125° , is yellow and on oxidation with nitric acid, forms phthalic acid.

β -Naphthoquinone decomposes about 120° . It crystallizes in red needles. It is odourless and non-volatile.

Anthraquinone, $C_{14}H_8O_2$, M.P. 273° , when treated with zinc dust and caustic soda, produces oxanthranol, $C_6H_4 \begin{smallmatrix} \text{CHOH} \\ \text{CO} \end{smallmatrix} C_6H_4$, which is red. The latter compound is easily oxidized on standing in air, anthraquinone being again formed.

Phenanthraquinone, M.P. 205° , is not reduced by sulphur dioxide. It dissolves in concentrated sulphuric acid to a dark green solution. When ignited with zinc dust, phenanthraquinone gives phenanthrene. The monoxime $C_{14}H_8O:NOH$ melts at 158° .

Chloranil (tetrachlorquinone), $C_6Cl_4O_2$, forms golden-yellow leaflets, with a smell reminiscent of Harris tweed. It sublimes undecomposed. With caustic potash it forms the purple potassium chloranilate, $C_6Cl_2(OK)_2O_2$. By the addition of acid to the latter, chloranilic acid, which crystallizes in reddish scales, is liberated.

XIX. CARBOHYDRATES.

THESE are solids, which, with the exception of starch, are soluble in water.

***d*-Glucose** (dextrose), $C_6H_{12}O_6 \cdot H_2O$, M.P. 86° . The anhydrous compound melts at 145° . Glucose reduces Fehling's solution. Glucosazone melts at $204-205^\circ$. Glucose pentabenzoate melts at 179° .

PREPARATION OF THE OSAZONE.—One gram of the sugar is dissolved in 5 cc. water, and 4 grams phenylhydrazine in 5 grams glacial acetic acid are added. The solution is heated on a water-bath for about ten minutes, when the osazone separates in yellow crystals. These are dried and the melting-point determined. This should be done quickly as the osazone decomposes to some extent in the neighbourhood of the melting-point.

Galactose, M.P. $163-164^\circ$. The osazone melts at $193-194^\circ$.

Fructose (laevulose), M.P. 95° . With phenylhydrazine it gives *d*-glucosazone. The pentabenzoate melts at $78-79^\circ$.

Cane Sugar, $C_{12}H_{22}O_{11}$, melting at 160° , does not reduce Fehling's solution. When boiled with dilute acids, glucose and fructose are produced. It forms no osazone. With benzoyl chloride it gives a hexabenzoate, melting at 109° .

Lactose (Milk sugar), $C_{12}H_{22}O_{11} \cdot H_2O$, becomes anhydrous at 140° , and melts with decomposition

at 205°. The osazone melts at 200°. Lactose reduces Fehling's solution.

Maltose, $C_{12}H_{22}O_{11}, H_2O$, loses water at 100°. When boiled with dilute acids, glucose is formed. The osazone melts at 190–191°.

Starch ($C_6H_{10}O_5$)_n, with iodine gives a blue colour, which disappears on heating. Starch is hydrolysed by heating with dilute acids, forming glucose. On boiling with water starch swells up and partially dissolves.

XX. GLUCOSIDES.

THESE, on hydrolysis with dilute acids or alkalies, give a sugar—generally glucose—and other substances.

Myronic Acid, $C_{10}H_{19}O_{10}NS_2$, usually occurs as the potassium salt. When boiled with barium hydroxide it is decomposed into glucose, allyl isothiocyanate (*see* p. 92) and potassium hydrogen sulphate.

Arbutin, $C_6H_4 \begin{matrix} \diagup OC_6H_{11}O_5 (1) \\ \diagdown OH (4) \end{matrix}$, is soluble in water.

When ferric chloride is added to the aqueous solution, the latter is turned deep blue. On hydrolysis with dilute sulphuric acid, glucose and hydroquinone are produced.

Salicin, $C_6H_4 \begin{matrix} \diagup OC_6H_{11}O_5 (1) \\ \diagdown CH_2OH (2) \end{matrix}$, melting at 201°, on

hydrolysis gives glucose and saligenin. Concentrated sulphuric acid produces a deep red colour. It is oxidized by chromic acid mixture to carbon dioxide, formic acid and salicyl aldehyde.

Amygdalin, $C_{20}H_{27}O_{11}N, 3H_2O$, loses water at 120° , melting at 200° and is hydrolysed by dilute acids into glucose, benzaldehyde and hydrocyanic acid..


Helicin, $C_6H_4 \begin{smallmatrix} \text{OC}_6\text{H}_{11}\text{O}_5 \\ \text{CHO} \end{smallmatrix}$, $\frac{1}{2}H_2O$, loses water at 100° and melts at 175° . On hydrolysis with dilute acids it gives glucose and salicyl aldehyde.

XXI. AMINES.


ALIPHATIC AMINES.

THE lower members are inflammable gases possessing an ammoniacal odour and are readily soluble in water, while the higher members are liquids. With acids they form salts which are soluble in water and in alcohol.


Nitrous Acid converts *primary* amines into the corresponding alcohols, with evolution of nitrogen—



Secondary amines are converted into yellow nitrosoamines—




Tertiary amines are unacted upon.



METHOD.—A small quantity of the amine is dissolved in dilute hydrochloric acid, care being taken that an excess of acid is present. To the cooled solution, a solution of sodium nitrite is added until free nitrous acid is present (starch iodine test).

Primary amines are also distinguished by giving the carbylamine test with chloroform and alcoholic potash—



METHOD.—About 0.1 gram of the amine is mixed with 3 drops chloroform and about 2 cc. alcoholic potash solution, and gently heated. A disgusting smell of carbylamine is evolved.

N.B.—The lower aliphatic amines are usually met with as salts.

Methylamine Hydrochloride, $CH_3NH_2 \cdot HCl$, forms deliquescent crystals.

Ethylamine, $C_2H_5NH_2$, B.P. 18–19°. The hydrochloride melts at 76–80°.

Diethylamine, $(C_2H_5)_2NH$, B.P. 56°. M.P. of hydrochloride 215–217°.

Triethylamine, $(C_2H_5)_3N$, B.P. 89°.

Propylamine, $(C_3H_7)NH_2$, B.P. 49°. The hydrochloride melts at 157–158°.

Isopropylamine, $(CH_3)_2CHNH_2$, B.P. 31–32°. The hydrochloride is deliquescent.

n-**Butylamine**, $C_4H_9NH_2$, B.P. 76°.

n-**Amylamine**, $C_5H_{11}NH_2$, B.P. 103°.

Allylamine, $CH_2 : CH \cdot CH_2NH_2$, B.P. 56°.

Ethylenediamine, $C_2H_4(NH_2)_2$, B.P. 116°. Nitrous acid converts it into ethylene oxide,

Tetramethylenediamine (putrescine), $C_4H_8(NH_2)_2$,
M.P. 27–28°.

Pentamethylene Diamine (cadaverine), $C_5H_{10}(NH_2)_2$.
B.P. 178–179°.

AROMATIC AMINES.

These are liquids or solids, and behave in many respects like the aliphatic amines. The *primary* compounds give the carbylamine reaction.

ACTION OF NITROUS ACID.—If the amino group is in the *nucleus* there are formed in cold acid solution diazonium salts, which on heating evolve nitrogen and produce phenols. If to a solution of a diazonium salt β -naphthol in caustic soda is added, a coloured precipitate of an azo-compound appears.

PREPARATION OF A DIAZONIUM SALT.—The amine is dissolved in a large excess of dilute hydrochloric acid and cooled in ice-water. Sodium nitrite solution is added in small quantities at a time, the mixture being well shaken after each addition, until, after standing for a few minutes, free nitrous acid can be detected by starch-iodide paper.

If the amino group is in the *side chain*, no diazonium salt is formed, and the amine behaves like the aliphatic compounds.

With *secondary* aromatic amines, nitroso-derivatives are formed, as in the case of aliphatic compounds.

Dialkylanilines with nitrous acid form para-nitroso compounds, the free bases of which are usually green, while the hydrochlorides are orange-coloured.

To completely identify an aromatic amine, its acetyl compound is prepared and its melting-point determined.

METHOD.—To 1 gram of the amine is added 1 gram acetic anhydride. The solid acetyl compound usually separates immediately. It is recrystallized from water or alcohol and its melting-point taken.

Aniline, $C_6H_5NH_2$, B.P. 182–183°. A chip of pine wood dipped first in hydrochloric acid and then in aniline, is turned yellow. Bleaching powder solution, when shaken with aniline becomes violet. On diazotizing and warming, aniline is converted into phenol. Acetanilide melts at 112°.

Methylaniline, $C_6H_5NHCH_3$, B.P. 193–194°, forms a nitroso-derivative, $C_6H_5NCH_3 \cdot NO$, M.P. 12–15°. The acetyl derivative melts at 101–102°.

Dimethylaniline, $C_6H_5N(CH_3)_2$, boils at 195°. The para-nitroso compound, which can be separated from its salts by means of sodium carbonate and extraction with ether, forms green crystals of metallic appearance, melting at 85°.

Ethylaniline, $C_6H_5NHC_2H_5$, B.P. 206°. The acetyl derivative melts at 54.5°.

Diethylaniline, $C_6H_5N(C_2H_5)_2$, B.P. 213–214°. *p*-Nitroso-compound M.P. 84°.

Diphenylamine, $(C_6H_5)_2NH$, melts at 54° and has a pleasant odour. The nitroso-derivative melts at 66.5°.

Triphenylamine, $(C_6H_5)_3N$, M.P. 127°. With concentrated sulphuric acid, there is produced a violet colouration, which changes to green.

o-Toluidine, $C_6H_4 \begin{smallmatrix} \text{CH}_3 \\ \text{NH}_2 \end{smallmatrix}$, boils at 197° . Ferric chloride precipitates a blue compound from its solution in hydrochloric acid. The acetyl compound melts at 110° .

m-Toluidine, B.P. 202° . The acetyl compound melts at 65° .

p-Toluidine, M.P. 45° . The melting point of the acetyl compound is 153° .


α -Naphthylamine, $C_{10}H_7NH_2$, M.P. 50° . On oxidation with chromic acid, α -naphthaquinone is produced.

β -Naphthylamine, M.P. $111-112^\circ$. Potassium permanganate oxidizes it to phthalic acid.


Benzylamine, $C_6H_5 \cdot CH_2NH_2$, B.P. $183-185^\circ$ is strongly alkaline.

o-Phenylenediamine, $C_6H_4 \begin{smallmatrix} \text{NH}_2 \\ \text{NH}_2 \end{smallmatrix}$, M.P. 102° . When ferric chloride is added to the hydrochloric acid solution, a dark red colouration is produced, and there quickly separate red needles of diamidophenazine $C_6H_4 \begin{smallmatrix} \text{N} \\ \text{N} \end{smallmatrix} > C_6H_4 (NH_2)_2$. The di-acetyl compound of *o*-phenylenediamine melts at $185-186^\circ$.

m-Phenylenediamine, M.P. 63° . Dilute nitrous acid produces a brown colouration owing to the formation



p-Phenylenediamine, M.P. 147°. When about 0.1 gram is dissolved in dilute hydrochloric acid and a little sulphuretted hydrogen water added along with a few drops of ferric chloride solution, and the mixture warmed, there appears a deep violet colouration (the so-called Lauth violet),



Benzidine, $\text{NH}_2\text{C}_6\text{H}_4\cdot\text{C}_6\text{H}_4\text{NH}_2$, melting at 122°, forms silvery leaflets, easily soluble in hot water and alcohol. The sulphate is insoluble in water.

HALOGEN AMINES.

***o*-Chloroaniline**, $\text{C}_6\text{H}_4\begin{smallmatrix} \text{Cl} \\ \text{NH}_2 \end{smallmatrix}$, B.P. 207°. The acetyl compound melts at 87-88°.


***m*-Chloroaniline**, B.P. 230°. M.P. of acetyl compound, 72.5°.

***p*-Chloroaniline**, M.P. 70°. M.P. of acetyl compound, 172.5°.

***o*-Bromoaniline**, $\text{C}_6\text{H}_4\begin{smallmatrix} \text{Br} \\ \text{NH}_2 \end{smallmatrix}$, M.P. 31°. M.P. of acetyl compound, 99°.

***m*-Bromoaniline**, M.P. 18°, B.P. 251°. M.P. of acetyl compound, 87.5°.

***p*-Bromoaniline**, M.P. 63°. M.P. of acetyl compound, 167-168°.



77.5°. M.P. of acetyl compound, 204°.


Sym. Tribromoaniline, M.P. 119–120°. M.P. of acetyl compound, 232°.

XXII. NITRO-COMPOUNDS.

ALIPHATIC NITRO-COMPOUNDS.


THESE are agreeably-smelling liquids, soluble in water. Many of them dissolve in alkalis.

ACTION OF NITROUS ACID.—With a *primary* aliphatic nitro-compound nitrous acid produces a red colouration due to the formation of a *nitrolic acid*.



The potassium salt is red.

With a *secondary* compound there is produced a dark blue colouration, a pseudo-nitrol being formed—



Tertiary compounds are unacted upon by nitrous acid.

METHOD.—A small quantity of the compound is shaken in a test-tube with sufficient caustic soda to form a clear solution. A few drops of a solution of sodium nitrite are added and finally dilute sulphuric

acid drop by drop. In the case of a primary compound excess of acid destroys the red colouration, which, however, is restored by addition of caustic soda.

REDUCTION.—Nitro-compounds are readily reduced to amines.

METHOD.—A few pieces of granulated zinc are placed on a test-tube and covered with caustic potash solution. Two cc. of the nitro-compound are introduced and the mixture warmed. The amine will be observed by its smell and its action on red litmus paper.

Nitromethane, CH_3NO_2 , B.P. 101° . Alcoholic caustic soda precipitates the sodium compound, $\text{CH}_2\text{:NO}_2\text{Na}$. Concentrated hydrochloric acid decomposes it with formation of formic acid.

Nitroethane, $\text{C}_2\text{H}_5\text{NO}_2$, B.P. $114\text{--}115^\circ$. Concentrated hydrochloric acid converts it into acetic acid.

Nitropropane, $\text{C}_3\text{H}_7\text{NO}_2$, B.P. $130\text{--}131^\circ$.

Tertiary Nitrobutane, $(\text{CH}_3)_3\text{CNO}_2$, is crystalline, melting at 24° . It is insoluble in alkalis.

AROMATIC NITRO-COMPOUNDS.

These are generally yellow oils or solids, insoluble in water, dilute hydrochloric acid or dilute caustic soda solution. Di-nitro and tri-nitro compounds impart a deep yellow colour to caustic soda solution.

They are reduced to amines by acid reducing agents.

METHOD.—A small quantity of the nitro-compound

is treated with zinc dust and hydrochloric acid, heat being applied if necessary. The mixture is diluted and rendered alkaline with caustic soda. It is then extracted with ether and the amine identified.

Nitrobenzene, $C_6H_5NO_2$, B.P. 209° , is a yellow oil with an odour of bitter almonds. On reduction it yields aniline.

***o*-Nitrotoluene**, $C_6H_4\begin{smallmatrix} CH_3 \\ \diagdown \\ NO_2 \end{smallmatrix}$, boils at 218° . It gives *o*-toluidine on reduction.

***m*-Nitrotoluene**, melts at 16° , and boils at $230-231^\circ$.

***p*-Nitrotoluene**, M.P. 54° .

***o*-Dinitrobenzene**, $C_6H_4\begin{smallmatrix} NO_2 \\ \diagdown \\ NO_2 \end{smallmatrix}$, M.P. $117-118^\circ$. On reduction it gives *o*-phenylenediamine.

***m*-Dinitrobenzene**, M.P. 91° .

***p*-Dinitrobenzene**, M.P. $171-172^\circ$.

***a*-Nitronaphthalene**, $C_{10}H_7NO_2$, melting at 61° , on reduction yields *a*-naphthylamine and with chromic acid is oxidized to nitrophthalic acid.

***β*-Nitronaphthalene**, M.P. 79° , gives on reduction *β*-naphthylamine.

NITROPHENOLS.


These compounds, on reduction with tin and hydrochloric acid, yield aminophenols.

***o*-Nitrophenol**, $C_6H_4\begin{smallmatrix} NO_2 \\ \diagdown \\ OH \end{smallmatrix}$, melting at 44° , has an intense yellow colour and a peculiar odour. It is

easily volatile in steam. The sodium salt gives a red solution in water.

m-Nitrophenol, M.P. 96° , forms sulphur-yellow crystals.

p-Nitrophenol, M.P. 114° , forms colourless needles which turn pink on exposure to air. The potassium salt is yellow. Phosphorus pentachloride converts the phenol into *p*-chloronitrobenzene, M.P. 83° .



yellow leaflets soluble in hot water. With aromatic hydrocarbons crystalline molecular compounds are obtained (see Naphthalene, p. 17). When picric acid is warmed with a concentrated solution of potassium cyanide, a dark red solution of isopurpuric acid is produced.

NITROKETONES.

o-Nitroacetophenone, $C_6H_4NO_2CO\cdot CH_3$, is an oil with a peculiar smell. It is oxidized by potassium permanganate to *o*-nitrobenzoic acid.

m-Nitroacetophenone forms needles melting at $80-81^{\circ}$. The oxime melts at 131° .

p-Nitroacetophenone, crystallizes in yellow prisms, melting at $80-81^{\circ}$.

NITROANILINES.

These are solids, yellow or orange in colour. They


have basic properties and are soluble in acids. On reduction with tin and hydrochloric acid the corresponding diamines are produced.

o-Nitroaniline, $C_6H_4 \begin{smallmatrix} \text{NO}_2 \\ \text{NH}_2 \end{smallmatrix}$, crystallizes in orange-

yellow needles, melting at 71° . When boiled with alkali it loses ammonia with production of *o*-nitrophenol. The acetyl derivative melts at 92° .

m-Nitroaniline, melting at 114° , forms yellow needles and is not acted on by alkali. The acetyl compound melts at 150° .

p-Nitroaniline, M.P. 147° , is decomposed by boiling alkali solution into ammonia and *p*-nitrophenol. The acetyl compound melts at 207° .



needles melting at 188° . When heated with alkali, alkali picrate is formed, and ammonia is evolved.

XXIII. NITROSO-COMPOUNDS.

Nitrosobenzene, $C_6H_5\text{NO}$, M.P. 68° , when fused or in solution has an intense greenish-blue colour.

p-Nitrosoaniline, $C_6H_4 \begin{smallmatrix} \text{NO} \\ \text{NH}_2 \end{smallmatrix}$, forms steel-blue


needles, melting at 174°. On boiling with caustic soda, it forms ammonia and *p*-nitrosophenol.

p-Nitrosophenol, $\text{C}_6\text{H}_4\text{<}\begin{smallmatrix}\text{NO} \\ \text{OH}\end{smallmatrix}$, M.P. 126°, is slightly

soluble in water, giving a light green solution. It is reduced by tin and hydrochloric acid to *p*-aminophenol. Concentrated nitric acid oxidizes it to *p*-nitrophenol.

Methyl-*p*-nitrosoaniline, $\text{C}_6\text{H}_4\text{<}\begin{smallmatrix}\text{NO} \\ \text{NHCH}_3\end{smallmatrix}$ ⁽¹⁾₍₄₎, forms

large crystals with metallic lustre, M.P. 118°. When boiled with caustic soda solution, it is decomposed into nitrosophenol and methylamine.



p-Nitrosodimethylaniline, $\text{C}_6\text{H}_4\text{<}\begin{smallmatrix}\text{NO} \\ \text{N}(\text{CH}_3)_2\end{smallmatrix}$, forms

green crystals, melting at 85°. The hydrochloride is yellow and is soluble with difficulty in water. Dilute caustic soda solution on warming decomposes the base into *p*-nitrosophenol and dimethylamine. It does not give Liebermann's nitroso reaction.


α-Nitrosonaphthol (*α*-naphthoquinone oxime), $\text{C}_{10}\text{H}_6(\text{OH})(\text{NO})$ 1 : 4, forms needles melting with decomposition at 193–194°.

β-Nitrosonaphthol (*β*-naphthoquinone oxime), $\text{C}_{10}\text{H}_6(\text{OH})(\text{NO})$ 1 : 2, crystallizes in yellowish-green needles, melting with decomposition at 152°. It dissolves in concentrated sulphuric acid with an intense red colour. When a neutral solution of the


sodium salt of β -nitroso-naphthol is added to a cobalt salt solution, a brown-red precipitate is formed.

XXIV. NITRILES AND ISONITRILES.

THE nitriles are liquids or solids of low melting-point, and have an agreeable odour. When boiled with strong mineral acids they are hydrolysed into the corresponding acids—



With nascent hydrogen they are reduced to amines.



HYDROLYSIS OF NITRILES.—Two cc. of the substance are boiled for some time with about 20 cc. concentrated hydrochloric acid. The liquid is then rendered alkaline with caustic soda and heated.

The residual liquid is tested for the fatty acid.


REDUCTION OF NITRILES.—One cc. of the substance is treated with 10 cc. dilute hydrochloric acid and a little zinc dust. The solution is rendered alkaline with caustic soda, and the amine identified.

Acetonitrile, $\text{CH}_3\cdot\text{CN}$. is a colourless liquid, boiling at $81-82^\circ$.

Benzonitrile, $\text{C}_6\text{H}_5\cdot\text{CN}$, B.P. 191° , is an oil with the odour of bitter almonds.

Isonitriles or **Carbylamines** are not usually met with unless in testing for primary amines. They are

liquids possessing a disgusting odour. On hydrolysis with acids they yield formic acid and an amine.



XXV. ISOCYANATES.

THESE are liquids, which, when digested with caustic potash yield primary amines and potassium carbonate, and with acids, a salt of a primary amine and carbon dioxide—



Methyl Isocyanate, B.P. 44°.




Ethyl Isocyanate, B.P. 60°.

Allyl Isocyanate, B.P. 82°.

Phenyl Isocyanate, B.P. 165–166°.


XXVI. UREAS AND UREIDES.

WHEN ureas and alkyl ureas are digested with caustic soda solution sodium carbonate remains, while from the former ammonia is evolved, from the latter amines. In the case of ureides, ammonia or amines may be given off, and the residue consists of sodium carbonate and the sodium salt of the acid of the ureide.



Urea, $\begin{array}{c} \text{NH}_2 \\ \diagup \text{CO} \\ \diagdown \text{NH}_2 \end{array}$, M.P. 132–133°, is a colourless solid.

When treated with nitrous acid it is decomposed into nitrogen and carbon dioxide—



With a solution of sodium hypobromite it gives the same decomposition products—



BIURET REACTION.—When urea is heated above its melting-point, ammonia is evolved and a residue of biuret remains—



When the biuret is dissolved in water and copper sulphate added and then caustic potash drop by drop, a violet colouration appears.

Methylurea, $\begin{array}{c} \text{NHCH}_3 \\ \diagup \text{CO} \\ \diagdown \text{NH}_2 \end{array}$, crystallizes in prisms,

melting at 102°. On heating for some time it decomposes into ammonia, methylamine and the dimethyl ester of cyanuric acid, which melts at 222°.

Ethylurea, $\begin{array}{c} \text{NHC}_2\text{H}_5 \\ \diagup \\ \text{CO} \\ \diagdown \\ \text{NH}_2 \end{array}$, melts at 92° . It decomposes on heating, into ammonia, ethylamine and the diethyl ester of cyanuric acid, M.P. 173° .


Parabanic Acid (Oxalylurea), $\begin{array}{c} \text{CO}-\text{NH} \\ | \quad \diagup \\ \text{CO}-\text{NH} \end{array} > \text{CO}$, on heating with caustic potash solution gives potassium oxalate, potassium carbonate and ammonia.

Barbituric Acid (Malonylurea), $\begin{array}{c} \text{NHCO} \\ \diagup \\ \text{CO} \\ \diagdown \\ \text{NHCO} \end{array} > \text{CH}_2$, on heating with caustic potash solution yields potassium malonate, potassium carbonate and ammonia.

Alloxan (Mesoxalylurea), $\begin{array}{c} \text{NHCO} \\ \diagup \\ \text{CO} \\ \diagdown \\ \text{NHCO} \end{array} > \text{CO}$, has an acid reaction. Ferrous sulphate gives a deep blue colour to the solution. When alloxan is boiled with dilute nitric acid, it is decomposed into parabanic acid and carbon dioxide.

Guanidine, $\begin{array}{c} \text{NH}_2 \\ \diagup \\ \text{CNH} \\ \diagdown \\ \text{NH}_2 \end{array}$, is a deliquescent strongly basic substance. The nitrate, M.P. 214° , dissolves with difficulty in water. When boiled with alkalis, guanidine is decomposed giving ammonia and alkali carbonate.

XXVII. URIC ACID GROUP.



alcohol, and practically insoluble in water.


1. When the acid is heated with soda lime ammonia is evolved.

2. **MUREXIDE TEST.**—A small quantity of the acid is moistened with concentrated nitric acid, and the mixture evaporated to dryness in a porcelain basin on a water-bath. There remains a reddish residue, which on addition of dilute ammonia solution changes to purple red, while alkali produces a violet colouration.

3. Fehling's solution is reduced by uric acid.

4. If a little of the acid be dissolved in a drop of caustic soda solution and this placed on filter paper which has been moistened with a solution of silver nitrate, a dark brown spot of metallic silver is immediately produced.


5. On dry distillation, uric acid decomposes without melting, giving off ammonia and hydrocyanic acid.



powder, soluble with difficulty in water, but very easily soluble in alkalis.

When a small quantity of xanthine is warmed with freshly-made chlorine water and a trace of nitric acid until evolution of gas ceases, the solution then carefully evaporated to dryness, and the solid exposed to ammonia gas, a rose-red colour is produced.

If xanthine be evaporated with nitric acid (Sp. Gr. 1.4), the residue is yellow. On addition of caustic potash it becomes yellowish red, and on warming violet-red.



is fairly soluble in water and in alcohol.

When evaporated with concentrated nitric acid it gives the murexide test (*see Uric Acid*).

When caffeine is evaporated with chlorine water, it leaves a purple red residue, which on strongly heating becomes yellow, but on addition of ammonia is changed to red.

XXVIII. HALOGEN COMPOUNDS.

ALIPHATIC.

THE alkyl halides are liquids which have a high specific gravity, and are practically insoluble in water. They are hydrolysed to the corresponding alcohols by warming with an excess of alkali solution.

Ethyl bromide, C_2H_5Br , B.P. 39° .

n-Propyl bromide, C_3H_7Br , B.P. 71° .

Isopropyl bromide, $CH_3\cdot CHBr\cdot CH_3$, B.P. $59-60^\circ$.

Methyl iodide, B.P. 43° .

Ethyl iodide, B.P. 72° .

Propyl iodide, B.P. 102° .

Isopropyl iodide, B.P. 89.5° .

Allyl chloride, $CH_2 : CH\cdot CH_2Cl$, B.P. 46° , has a leek-like odour.

Allyl bromide, B.P. $70-71^\circ$.

Allyl iodide, B.P. $102-103^\circ$.

Methylene iodide, CH_2I_2 , boils at 180° with partial decomposition.

Ethylene chloride, $CH_2Cl\cdot CH_2Cl$, B.P. 84° .

Ethylene bromide, B.P. $131-132^\circ$.

Ethylene iodide, melts at $81-82^\circ$.

Chloroform, $CHCl_3$, B.P. 61.5° , is a colourless non-inflammable liquid. When boiled with alcoholic caustic potash, potassium formate and potassium chloride are formed—



The carbylamine test may also be applied (*see* page 11).

Bromoform, CHBr_3 , B.P. 151° .

Iodoform, CHI_3 , M.P. $119\text{--}120^\circ$, crystallizes in brilliant yellow leaflets of a characteristic odour.

Carbon tetrachloride, CCl_4 , B.P. $76\text{--}77^\circ$, is a pleasant smelling liquid. When treated with zinc and hydrochloric acid, it is reduced to chloroform. On heating with alcoholic potash it is decomposed into potassium carbonate and potassium chloride.



AROMATIC HALOGEN COMPOUNDS

are divided into two classes :—

1. True aromatic halogen compounds, i.e. those with the halogen attached to the benzene nucleus.
2. Those with the halogen in the side chain.

The members of the first class are colourless liquids or solids, with a faint, agreeable odour. They are insoluble in water, but readily soluble in the other common solvents. They easily form nitro-derivatives. Caustic alkali does not readily remove the halogen from these compounds.

Monochlorobenzene, $\text{C}_6\text{H}_5\text{Cl}$, B.P. 132° .

***o*-Dichlorobenzene**, $\text{C}_6\text{H}_4\text{Cl}_2$, B.P. 179° .

***m*-Dichlorobenzene**, B.P. 172° .

***p*-Dichlorobenzene**, M.P. 53° .

Monobromobenzene, $\text{C}_6\text{H}_5\text{Br}$, B.P. $156\text{--}157^\circ$.

***o*-Dibromobenzene**, $\text{C}_6\text{H}_4\text{Br}_2$, B.P. 224° .

***m*-Dibromobenzene**, B.P. 219° .

p-Dibromobenzene, M.P. 89°.

Iodobenzene, C_6H_5I , B.P. 188°.

o-Chlorotoluene, $C_6H_4 \begin{smallmatrix} \text{CH}_3 \\ \text{Cl} \end{smallmatrix}$, B.P. 156°.

m-Chlorotoluene, B.P. 150°.

p-Chlorotoluene, B.P. 163°.

o-Bromotoluene, $C_6H_4 \begin{smallmatrix} \text{CH}_3 \\ \text{Br} \end{smallmatrix}$, B.P. 181°.

m-Bromotoluene, B.P. 183–184°.


p-Bromotoluene, M.P. 28–29°.

α -Chloronaphthalene, $C_{10}H_7Cl$, B.P. 263°.


β -Chloronaphthalene, M.P. 56°, B.P. 265°.

The compounds with halogen in the side chain behave like aliphatic halogen compounds.


Benzyl chloride, $C_6H_5 \cdot CH_2Cl$, B.P. 175°, when boiled with a solution of copper nitrate, benzaldehyde is formed—



Benzal chloride (benzylidene chloride), $C_6H_5 \cdot CHCl_2$, B.P. 206°, forms benzaldehyde, when heated with milk of lime—








Benzotrichloride, $C_6H_5 \cdot CCl_3$, B.P. 213°, on heating with milk of lime, gives calcium benzoate—



XXIX. AZO COMPOUNDS.

THESE compounds are coloured solids, which on reduction produce amino compounds.



METHOD OF REDUCTION.—About 3 grams of the compound are warmed with zinc dust and water until the colour has disappeared. The mixture is then extracted with ether and the ethereal solution evaporated to dryness. The residue may be :—

1. A mixture of two amines.
2. A mixture of an amine and an aminophenol.
3. A mixture of an amine and an aminosulphonic acid.

MIXTURE OF TWO AMINES.—The residue of liquid is distilled and the first and last fractions treated with acetic anhydride. The acetyl compounds thus formed are then identified by their melting points.

MIXTURE OF AN AMINE AND AN AMINOPHENOL.—The aminophenol is extracted from the mixture with alkali.

MIXTURE OF AN AMINE AND AN AMINOSULPHONIC ACID.—The mixture is neutralized with alkali, the amine extracted with ether, and identified. The aminosulphonic acid is liberated by adding hydrochloric acid to the alkali salt.

Azobenzene, $C_6H_5N : NC_6H_5$, M.P. 68° , is easily soluble in alcohol. When treated with tin and hydrochloric acid, benzidine is produced.

p-Aminoazobenzene, $C_6H_5N : NC_6H_4NH_2$, forms yellow leaflets melting at $125-126^\circ$. It is oxidized by sulphuric acid and manganese dioxide with formation of quinone. The hydrochloride crystallizes in steel-blue needles. The acetyl derivative melts at 142° .

Diaminoazobenzene, $C_6H_5N : NC_6H_3 \begin{matrix} \swarrow NH_2 (1) \\ \searrow NH_2 (3) \end{matrix}$

forms yellow needles, M.P. 117° . On reduction it yields aniline and triaminobenzene, M.P. 132° . The hydrochloride of diaminoazobenzene is known as **chrysoidine**.

Triaminoazobenzene, $NH_2C_6H_4N : NC_6H_3 \begin{matrix} \swarrow NH_2 (2) \\ \searrow NH_2 (4) \end{matrix}$,

M.P. 143° .

Helianthine, $SO_3HC_6H_4N : NC_6H_4N(CH_3)_2$, forms glistening violet leaflets. On reduction it yields sulphanilic acid and p-amino-dimethylaniline, M.P. 41° . The sodium salt is known as **Methyl Orange**.

The following, which are related to the azo-compounds, should also be mentioned.

Hydrazobenzene, $C_6H_5NH \cdot NHC_6H_5$, M.P. 131° , forms colourless leaflets, which are insoluble in water.

If its alcoholic solution be exposed to air, azobenzene is produced. Powerful reducing agents decompose it into aniline.

Azoxybenzene, $C_6H_5N - NC_6H_5$, crystallizes in long



yellow needles, M.P. 36° . It is insoluble in water. On distillation it decomposes into azobenzene and aniline.

Diazoaminobenzene, $C_6H_5N : N \cdot NHC_6H_5$, melts at 96° and explodes at higher temperatures.

XXX. PYRIDINE AND QUINOLINE GROUP.

Pyridine, C_5H_5N , B.P. 116° , has a strong characteristic odour, and is miscible in all proportions with water, alcohol and ether. It forms salts with acids. On reduction with sodium and alcohol, piperidine is produced.

Piperidine, $C_5H_{11}N$, B.P. 106° , is a colourless liquid with a characteristic odour. It is miscible with water, alcohol and ether. It is a secondary amine.

Quinoline, C_9H_7N , B.P. 239° , is sparingly soluble in water. The hydrochloride is crystalline. With methyl iodide quinoline forms a yellow crystalline compound melting at 72° .

XXXI. ALKALOIDS.

THE alkaloids are nearly all solids, easily soluble in alcohol, soluble with difficulty in ether, chloroform and benzene, and sparingly soluble or insoluble in water. Most are optically active, generally laevorotatory. They dissolve readily in dilute acids with formation of salts, from which the alkaloid is precipitated on addition of an alkali. Morphine, however, is soluble in excess of sodium hydroxide.

The following reactions are given by almost all the alkaloids :—

1. When an alkaloid is heated in a dry test-tube, decomposition takes place with production of a smell like burning feathers.

2. A solution of iodine in potassium iodide produces a brown flocculent precipitate in solutions of salts of the alkaloids, best when acidified with dilute sulphuric acid.


3. Phosphomolybdic acid¹ produces a yellow precipitate in solutions of all the alkaloids. The precipitate is soluble in alkalies and alkaline carbonates.

4. Potassium mercuric iodide² produces a white or yellowish white precipitate in solutions of salts of alkaloids. The precipitates are insoluble in dilute hydrochloric acid.

5. Aqueous solutions of tannic acid and of picric acid precipitate all or nearly all the alkaloids from solutions of their salts.

¹ See Appendix.

² See Appendix.



less oil with an unpleasant smell, B.P. 247°.

Picric acid produces an amorphous precipitate which readily crystallizes in small yellow needles melting at 218°.

Quinine, when anhydrous melts at 177°. The normal salts are sparingly soluble, the acid salts very soluble.

Solutions of quinine salts, when acidified with sulphuric acid show a fine blue fluorescence.

Concentrated sulphuric acid dissolves the alkaloid to a colourless solution, which on heating turns brown and yellow.

When a solution of a quinine salt is mixed with one-fifth of its volume of chlorine water and then with an excess of ammonia, an emerald green colour is produced. If a little potassium ferrocyanide solution be added after the chlorine water and then a few drops of ammonia, the solution becomes deep red in colour.

Cinchonine, M.P. 250°.

When chlorine water and an excess of ammonia are added to a solution of a cinchonine salt, a yellowish white precipitate is formed.

When potassium ferrocyanide is added to a neutral or slightly acid solution of a cinchonine salt, a yellowish white flocculent precipitate is produced.

When chlorine water is added drop by drop to a solution of a brucine salt, there is produced a red colouration which is destroyed by excess of chlorine.

Strychnine, M.P. 284°.

If a crystal of potassium dichromate be stirred in a solution of the alkaloid in concentrated sulphuric acid, there appears a fine blue colour, which changes successively to violet, red and reddish yellow.

Potassium ferrieyanide and also potassium chromate produce yellow crystalline precipitates in neutral and fairly concentrated solutions of strychnine salts.

Brucine, when anhydrous melts at 178°.

The solution in concentrated sulphuric acid has at first a rose-red colour, which changes to yellow.

Concentrated nitric acid dissolves brucine and its salts giving an intensely red solution, which, when warmed, turns yellow. If stannous chloride be added to this yellow solution, an intense violet colouration is produced.

Morphine is soluble in dilute acids and in alkalis. Its salts dissolve easily in water and alcohol.

The addition of concentrated nitric acid to the solid alkaloid or a salt causes a yellowish red colour to appear.

When ammonium molybdate in concentrated

sulphuric acid (0.1 gram in a cc.) is placed in a porcelain dish and a particle of morphine added and crushed with a glass rod, a deep violet colour appears, which slowly changes to blue.

If a few drops of a dilute neutral solution of ferric chloride be added to a concentrated solution of a morphine salt, a dark blue colouration is produced. The colour is destroyed by acids.

Narcotine, M.P. 176° .


Cold concentrated sulphuric acid dissolves the alkaloid, giving a yellow solution. If this be carefully heated in a porcelain dish, the colour changes from yellow through orange to red, while a bluish violet colour appears at the edges. If the heating be continued until the acid begins to evaporate, the solution acquires a reddish violet colour.

If to a solution of narcotine in concentrated sulphuric acid there be added 10 to 20 drops sulphuric acid containing a trace of nitric acid, the liquid turns brown and then quickly red.

XXXII. SULPHUR COMPOUNDS.

(OTHER THAN SULPHONIC ACIDS AND DERIVATIVES.)

Carbon Disulphide, CS_2 , is a highly refractive colourless liquid boiling at 47° . When it is added to alcoholic potash, potassium xanthate is precipitated in yellow needles—



MERCAPTANS OR THIO-ALCOHOLS.

These are colourless liquids with a disagreeable garlic-like odour, and are insoluble in water.

1. Oxidation with nitric acid produces sulphonic acids which can be identified by conversion into their amides (*see* page 48).

2. When a mercaptan is added to a few cc. of an alcoholic solution of mercuric chloride, a white precipitate is formed.

3. When sodium is added to a mercaptan in ethereal solution, hydrogen is evolved and the sodium salt crystallizes in white needles.

Ethyl mercaptan, $\text{C}_2\text{H}_5\text{SH}$, B.P. 36° .

n-Propyl mercaptan, $\text{C}_3\text{H}_7\text{SH}$, B.P. 68° .


Isopropyl mercaptan, $(\text{CH}_3)_2\text{CHSH}$, B.P. 59° .

THIOETHERS OR ALKYL SULPHIDES.

The thioethers are colourless liquids with a disagreeable odour, insoluble in water, but easily soluble in alcohol and ether. They are characterized by their additive powers; for example, they unite with alkyl iodides to form sulphine iodides.

One cc. ethyl iodide is added to 1 cc. of the thioether and the mixture warmed. After adding about a gram of moist silver oxide, the mixture is

shaken with water, again warmed and filtered. The filtrate will be strongly alkaline owing to the presence of trialkyl sulphoxide.



Methyl sulphide, $(CH_3)_2S$, B.P. 37.5° .

Ethyl sulphide, $(C_2H_5)_2S$, B.P. 92° .

***n*-Propyl Sulphide**, $(C_3H_7)_2S$, B.P. $130-135^\circ$.

Allyl Sulphide, $(CH_2\cdot CH : CH)_2S$, B.P. $139-140^\circ$, forms a crystalline precipitate with alcoholic mercuric chloride.

Thiophene, C_4H_4S , B.P. 84° .

When a crystal of isatin is dissolved in 2 cc. conc. sulphuric acid, and to this solution a few drops of thiophene are added, a deep blue colour is produced.

COMPOUNDS CONTAINING NITROGEN AND SULPHUR.

MUSTARD OILS OR ISOTHIOCYANATES.

These are liquids with a very penetrating odour. They are insoluble in water. Their boiling points are lower than those of the corresponding thio-cyanates.


1. When heated to 100° with concentrated hydrochloric acid under a reflux condenser, they are decomposed into amines, sulphuretted hydrogen and carbon dioxide—



2. Reduction by means of zinc and hydrochloric acid produces an amine and thioformaldehyde, which has an odour of onions—



3. One cc. of the mustard oil is warmed with one cc. aniline for a few minutes. On cooling, a thiourea crystallizes out, and its melting point is determined (*see* p. 93).



4. If 1 cc. of the mustard oil be heated with one gram yellow mercuric oxide, an isocyanate and a black precipitate of mercuric sulphide are formed.

Methyl mustard oil, $CSNCH_3$, M.P. 35° .


Ethyl mustard oil, $CSNC_2H_5$, B.P. $131-132^\circ$.




Allyl mustard oil, $CSNC_3H_5$, B.P. 150.7° .

Phenyl mustard oil, $CSNC_6H_5$, B.P. 222° , when reduced with zinc dust gives benzonitrile.








THIOUREAS.




When a thiourea is heated with concentrated hydrochloric acid under a reflux condenser for some time and then distilled, the distillate contains a mustard oil. If the residue, which contains a guanidine derivative, be heated with concentrated caustic soda solution, an amine and a carbonate are produced.






alkalies, hydrochloric acid or sulphuric acid, is decomposed.





THIOCYANATES.

1. When a thiocyanate is reduced with zinc dust and hydrochloric acid, a mercaptan and hydrocyanic acid¹ are produced—



2. Boiling nitric acid oxidizes thiocyanates to alkyl sulphonic acids.

Methyl thiocyanate, CNSCH_3 , B.P. 133°.

Ethyl thiocyanate, CNSC_2H_5 , B.P. 142°.

Allyl thiocyanate, CNSC_3H_5 , B.P. 161°, rapidly changes on boiling to the isomeric mustard oil.

XXXIII. TERPENES AND ALLIED COMPOUNDS.

THESE compounds, although possessing many of the properties of compounds already considered, have distinctive characteristics, thus necessitating

¹ Since hydrocyanic acid is a poison, great care must be taken in carrying out this reaction.

their separate classification. They are neither aliphatic nor aromatic, and are complex in structure. Nevertheless, they yield characteristic derivatives which as a general rule can be easily identified. They are all highly inflammable, possess characteristic odours and are insoluble in water, but are readily soluble in most organic solvents.

HYDROCARBONS.

Dipentene, $C_{10}H_{16}$, B.P. 175–176°. The *dihydrochloride*, $C_{10}H_{16} \cdot 2HCl$, is prepared by passing dry hydrochloric acid over a few cc. dipentene for about one hour. The dihydrochloride separates on pouring the mixture on ice. It is dried on a porous plate and recrystallized from alcohol. M.P. 48–50°.

Dipentene tetrabromide, $C_{10}H_{16}Br_4$, can be made by adding bromine to a chloroform solution of dipentene, cooled to -10° by salt and ice. The chloroform is removed by blowing a current of dry air through the solution. The tetrabromide on recrystallization from ethyl acetate, melts at 125° .

The active forms of dipentene are the **limonenes**. The active tetrabromides melt at 104° .

Pinene, $C_{10}H_{16}$, B.P. 155–156°.

The nitrosochloride, $C_{10}H_{16}NOCl$, melts at 115° . It is prepared by adding 1.5 cc. 33 per cent. hydrochloric acid to a mixture (cooled in ice) of 5 grams pinene, 5 grams glacial acetic acid, and 5 grams ethyl nitrite. After a short time the nitrosochloride

separates in large crystals. It is filtered off at the pump and washed with alcohol.

Camphene, $C_{10}H_{16}$, melts at 51° .

When heated with glacial acetic acid and a little sulphuric acid, camphene forms isobornyl acetate. On adding water the ester separates as an oil. It is hydrolysed with alcoholic caustic potash, the alcohol removed, and isborneol is precipitated on addition of water. Isborneol melts about 208° in a closed tube.

Menthene, $C_{10}H_{18}$, B.P. $167-168^{\circ}$. The nitroso-chloride melts at 127° .

ALCOHOLS.

Terpineol, $C_{10}H_{17}OH$, melts at 35° . It forms a nitrosochloride melting at $112-113^{\circ}$.

Borneol, $C_{10}H_{17}OH$, M.P. $203-204^{\circ}$, and **Iso-borneol**, $C_{10}H_{17}OH$, M.P. 208° , yield derivatives of very similar melting and boiling points. The phenylurethanes of both melt at $138-139^{\circ}$.

PREPARATION OF A PHENYLURETHANE.—Molecular quantities of the alcohol and phenyl isocyanate¹ are mixed and heated rapidly to boiling. The mixture is well shaken and allowed to stand, the flask being closed with a calcium chloride tube. The unchanged phenyl isocyanate is extracted with benzene, and the residual urethane, after removal of benzene, is washed with cold water and recrystallized from ethyl acetate or a mixture of

¹ This preparation should be carried out in a draught-cupboard.

ether and petroleum ether. (Solvents containing "hydroxyl" should be avoided.)

When isoborneol is heated for some time with ethyl alcohol and concentrated sulphuric acid, isobornyl ethyl ether (B.P. 203–204°) is produced. Bornyl ethyl ether is not formed in this way.

Menthol, $C_{10}H_{18}OH$, M.P. 42°. The phenylurethane forms needles, melting at 111°.

KETONES.

Carvone, $C_{10}H_{14}O$, B.P. 223–224°. On heating with a little acid it is converted into carvacrol.

Camphor, $C_{10}H_{16}O$, M.P. 177–178°. The oxime melts at 120°; the semicarbazone at 236°.

Menthone, $C_{10}H_{18}O$, B.P. 208°, has a peppermint-like odour. The semicarbazone melts at 184°.

XXXIV. ALBUMINS AND PROTEIDS.

THESE are compounds of very complicated structure, and, with a few exceptions, do not crystallize. They are insoluble in the common organic solvents, but dissolve in aqueous solutions of acids and alkalies.

The following are general tests for the albumins :—

I. **MILLON'S REAGENT** (a solution of mercuric nitrate containing nitrous acid) gives an intense red colouration on heating.

II. XANTHO-PROTEIN REACTION. — By heating an albumin with fairly concentrated nitric acid, there separates a yellow flocculent precipitate of xanthoproteic acid, which dissolves in alkali giving an orange red solution.

III. BIURET REACTION. — When caustic potash is added to albumin, and then a very dilute solution of copper sulphate drop by drop, a fine violet-red colouration appears.


Egg-albumin is soluble in water and is precipitated from its solutions by metaphosphoric acid and by acetic acid. The precipitate is soluble in excess of the latter. Egg-albumin treated with concentrated sulphuric acid and sugar solution gives a red solution which changes to violet-red.

Casein contains phosphorus. An alkaline solution of casein dissolves cupric hydrate, giving a violet colour.

Gelatine contains sulphur. It is precipitated from its solutions by tannin.

Appendix.

MELTING-POINT APPARATUS.—The sketch shows an apparatus which is very convenient and gives very



accurate determinations.

SPECIAL REAGENTS.

NEUTRAL FERRIC CHLORIDE SOLUTION.—To some of the ferric chloride solution provided in the laboratory, dilute ammonium hydroxide or ammonium carbonate is added drop by drop until a precipitate just begins to form.

SCHIFF'S REAGENT.—Sulphurous acid is added to a dilute solution of rosaniline hydrochloride (magenta) until the colour just disappears.

FEHLING'S SOLUTION.—(1) 35 grams copper sulphate are dissolved in water and diluted to 1 litre. (2) 173 grams sodium potassium tartrate (Rochelle salt) and 70 grams caustic soda are dissolved in water and diluted to 1 litre.

These solutions are kept in separate bottles. Equal volumes of each are mixed immediately before making a test.

SOLUTION OF IODINE IN POTASSIUM IODIDE.—5 grams of iodine are dissolved in water containing 10 grams of potassium iodide and the solution made up to 1 litre.

HYDRIODIC ACID.—To obtain the constant boiling solution (Sp. Gr. 1.7) hydriodic acid of Sp. Gr. 2.00 is diluted with an equal volume of water and distilled over a small quantity of red phosphorus. The fraction boiling at 125°–126° is collected separately.

SODIUM BISULPHITE (saturated solution).—Sulphur dioxide is passed into a saturated solution of sodium carbonate for some time. Solid carbonate is added, from time to time, until no more dissolves. The final solution should smell strongly of sulphur dioxide.

PHOSPHOMOLYBDIC ACID.—To a solution of 75 grams ammonium molybdate in 500 cc. nitric acid (Sp. Gr. 1.2) and 500 cc. water, sodium phosphate solution is added until there is no further precipitate. This is then filtered off, well washed and finally warmed with sodium carbonate solution until completely dissolved. The solution is evaporated to dryness, and the residue ignited. The product is washed with water and dissolved in a considerable excess of nitric acid.

POTASSIUM MERCURIC IODIDE.—13 grams mercuric chloride and 50 grams potassium iodide are dissolved in water and the mixture made up to 1 litre.

MILLON'S REAGENT.—A small quantity of mercury is dissolved in twice its weight of concentrated nitric acid in the cold and twice its volume of water is added.

INDEX

	PAGE		PAGE
A.		Aldehydes	29
Acetyl compounds of		Aldehydes substituted	33
amines, preparation of	65	Alkaloids	86
Acid amides	53	Alkyl sulphides	90
Acid anhydrides	52	Amines, aliphatic	62
Acid anilides	55	Amines, aromatic	64
Acid halides	53	Amines, halogen	67
Acid imides	55	Amino group, detection	
Acids, aliphatic mono-		of	11
basic saturated	38	Aminophenols	28
Acids, aliphatic poly-		Azo compounds	83
basic saturated	39	Azo group, detection of	14
Acids, aliphatic unsatu-			
rated	40	B.	
Acids, amino	46	Biuret reaction	76
Acids, aromatic	41	C.	
Acids, aromatic sulphonie	48	Carbohydrates	60
Acids, halogen substi-		Chromic acid oxidation	
tuted	45	mixture.	15
Acids, hydroxy	43	D.	
Acids, nitro	46	Diazonium salt, prepara-	
Albumins	97	tion of	64
Alcohol phenols	27	E.	
Alcohols, polyhydric	21	Esters	56
Alcohols, primary mono-		Ethers	22
hydric	19	Ethoxy group, detection	
Alcohols, secondary mono-		of	9
hydric	21		
Alcohols, tertiary mono-			
hydric	21		
Aldehyde ethers	36		
Aldehyde group, detec-			
tion of	9		

	PAGE		PAGE
F		M.	
Fehling's solution, preparation of	99	Melting-point apparatus	99
Ferric chloride, neutral solution.	99	Mendius' reaction	13, 74
		Mercaptans	90
G.		Methoxy group, detection of	9
Glucosides	61	Millon's reagent, preparation of	100
		Mustard oils	91
H.		N.	
Halogen compounds, aliphatic	80	Nitriles	74
Halogen compounds, aromatic	81	Nitriles, detection of	13
Halogen phenols.	27	Nitro compounds, aliphatic	68
Hydriodic acid, constant boiling	100	Nitro compounds, aromatic	69
Hydrocarbons, aliphatic	14	Nitro group, detection of	3
Hydrocarbons, aromatic	15	Nitroanilines	71
Hydroxyl group, detection of	8	Nitro-ketones	71
		Nitrophenols	70
		Nitroso compounds	72
		Nitroschloride, preparation of a	95
I.		O.	
Imino group, detection of	12	Oxidation with chromic anhydride	18
Iodine in potassium iodide.	100	Oximes, preparation of	29
Isocyanates	75		
Isonitriles.	74	P.	
Isonitriles, detection of	13	Phenols	23
Isothiocyantes	91	Phenols, alcohol	27
		Phenols, halogen	27
K.		Phenylhydrazones, preparation of	30
Ketone group, detection of	11	Phenylurethanes, preparation of	96
Ketones	36	Phosphomolybdic acid, preparation of	100
Ketones, nitro-	71	Picrates, preparation of	17
		Piperidine	85
L.			
Liebermann's test	12		

INDEX

103

	PAGE		PAGE
Potassium mercuric iodide, preparation of	100	Sulphur compounds.	89
Proteids	97		
Pyridine	85		
		T.	
Q.		Terpenes	94
Quinoline	85	Tests for elements	4
Quinones	58	Tests, preliminary	1
		Thioalcohols	90
S.		Thiocyanates.	94
Schiff's reaction	10	Thio-ethers	90
Schiff's reagent, preparation of	99	Thiophene	91
Schotten-Baumann reaction	8	Thioureas.	92
Semicarbazones, preparation of	30	Tollen's reaction	9
Sodium bisulphite, preparation of	100		
Sulphonamides, preparation of	48	U.	
		Ureas	75
		Ureides	75
		Unsaturation, tests for	4
		Uric acid group.	78

1

D. VAN NOSTRAND COMPANY

23 MURRAY AND 27 WARREN STREETS

NEW YORK


SHORT-TITLE CATALOG

OF

Publications and Importations

OF

**SCIENTIFIC AND ENGINEERING
BOOKS**



This list includes the technical publications of the following English publishers :

**SCOTT, GREENWOOD & CO. CROSBY LOCKWOOD & SON
CONSTABLE & COMPANY, Ltd. TECHNICAL PUBLISHING CO.
ELECTRICIAN PRINTING & PUBLISHING CO.,**

for whom D. Van Nostrand Company are American agents.

Descriptive Circulars sent on request.

SHORT-TITLE CATALOG

OF THE

Publications and Importations

OF

D. VAN NOSTRAND COMPANY
23 MURRAY AND 27 WARREN STREETS, N. Y.

Prices marked with an asterisk (*) are NET
 All bindings are in cloth unless otherwise noted

A B C Code. (See Clausen-Thue.)	
Abbott, A. V. The Electrical Transmission of Energy.....	8vo, *\$5 00
— A Treatise on Fuel. (Science Series No. 9.).....	16mo, 0 50
— Testing Machines. (Science Series No. 74.).....	16mo, 0 50
Adam, P. Practical Bookbinding. Trans. by T. E. Maw.....	12mo, *2 50
Adams, J. W. Sewers and Drains for Populous Districts....	8vo, 2 50
Addyman, F. T. Practical X-Ray Work.....	8vo, *4 00
A1 Code. (See Clausen-Thue.)	
Aikman, C. M. Manures and the Principles of Manuring. . .	8vo, 2 50
Alexander, J. H. Elementary Electrical Engineering.....	12mo, 2 00
— Universal Dictionary of Weights and Measures.....	8vo, 3 50
Allan, W. Strength of Beams under Transverse Loads. (Science Series No. 19.).....	16mo, 0 50
— Theory of Arches. (Science Series No. 11.).....	16mo, 0 50
Allen, H. Modern Power Gas Producer Practice and Applica- tions.....	12mo, *2 50
— Gas and Oil Engines.....	8vo, *4 50
Anderson, F. A. Boiler Feed Water.....	8vo, *2 50
Anderson, Cap. G. L. Handbook for the Use of Electricians .	8vo, 3 00
Anderson, J. W. Prospector's Handbook.....	12mo, 1 50

Andes, L. Vegetable Fats and Oils.....	8vo,	*4 00
— Animal Fats and Oils. Trans. by C. Salter.....	8vo,	*4 00
— Drying Oils, Boiled Oil, and Solid and Liquid Driers....	8vo,	*5 00
— Iron Corrosion, Anti-fouling and Anti-corrosive Paints. Trans. by C. Salter.....	8vo,	*4 00
— Oil Colors and Printers' Ink. Trans. by A. Morris and H. Robson.....	8vo,	*2 50
— Treatment of Paper for Special Purposes. Trans. by C. Salter.....	12mo,	*2 50
Annual Reports on the Progress of Chemistry.		
Vol. I. (1904).....	8vo,	*2 00
Vol. II. (1905).....	8vo,	*2 00
Vol. III. (1906).....	8vo,	*2 00
Vol. IV. (1907).....	8vo,	*2 00
Vol. V. (1908).....	8vo,	*2 00
Vol. VI. (1909).....	8vo,	*2 00
Argand, M. Imaginary Quantities. Translated from the French by A. S. Hardy. (Science Series No. 52.).....	16mo,	0 50
Armstrong, R., and Idell, F. E. Chimneys for Furnaces and Steam Boilers. (Science Series No. 1.).....	16mo,	0 50
Arnold, E. Armature Windings of Direct Current Dynamos. Trans. by F. B. DeGress.....	8vo,	*2 00
Ashe, S. W., and Keiley, J. D. Electric Railways. Theoretically and Practically Treated. Vol. I. Rolling Stock	12mo,	*2 50
Ashe, S. W. Electric Railways. Vol. II. Engineering Preliminaries and Direct Current Sub-Stations.....	12mo,	*2 50
— Electricity: Experimentally and Practically Applied.	12mo,	*2 00
Atkinson, A. A. Electrical and Magnetic Calculations.....	8vo,	*1 50
Atkinson, J. J. Friction of Air in Mines. (Science Series No. 14.).....	16mo,	0 50
Atkinson, J. J., and Williams, E. H., Jr. Gases Met with in Coal Mines. (Science Series No. 13.).....	16mo,	0 50
Atkinson, P. The Elements of Electric Lighting.....	12mo,	1 50
— The Elements of Dynamic Electricity and Magnetism. 12mo,		2 00
— Power Transmitted by Electricity.	12mo,	2 00
Auchincloss, W. S. Link and Valve Motions Simplified....	8vo,	*1 50
Ayrton, H. The Electric Arc.....	8vo,	*5 00

4 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Bacon, F. W. Treatise on the Richards Steam-Engine Indicator.....	12mo,	1 00
Bailes, G. M. Modern Mining Practice. Five Volumes. 8vo, each,		3 50
Bailey, R. D. The Brewers' Analyst.....	8vo,	*5 00
Baker, A. L. Quaternions.....	12mo,	*1 25
Baker, Benj. Pressure of Earthwork. (Science Series No. 56.)		
	16mo,	
Baker, I. O. Levelling. (Science Series No. 91.).....	16mo,	0 50
Baker, M. N. Potable Water. (Science Series).....	16mo,	0 50
— Sewerage and Sewage Purification. (Science Series No. 18.)		
	16mo,	0 50
Baker, T. T. Telegraphic Transmission of Photographs.....	12mo,	*1 25
Bale, G. R. Modern Iron Foundry Practice. Two Volumes.		
	12mo.	
Vol. I. Foundry Equipment, Material Used.....		*2 50
Vol. II. Machine Moulding and Moulding Machines.....		*1 50
Bale, M. P. Pumps and Pumping.....	12mo,	1 50
Ball, R. S. Popular Guide to the Heavens.....	8vo,	*4 50
— Natural Sources of Power. (Westminster Series).....	8vo,	*2 00
Ball, W. V. Law Affecting Engineers.....	8vo,	*3 50
Bankson, Lloyd. Slide Valve Diagrams. (Science Series No. 108.).....	16mo,	0 50
Barba, J. Use of Steel for Constructive Purposes.....	12mo,	1 00
Barker, A. Textiles and Their Manufacture. (Westminster Series).....	8vo,	2 00
Barker, A. H. Graphic Methods of Engine Design.....	12mo,	*1 50
Barnard, F. A. P. Report on Machinery and Processes of the Industrial Arts and Apparatus of the Exact Sciences at the Paris Universal Exposition, 1867.....	8vo,	5 00
Barnard, J. H. The Naval Militiaman's Guide.....	16mo, leather,	1 25
Barnard, Major J. G. Rotary Motion. (Science Series No. 90.)		
	16mo,	0 50
Barrus, G. H. Boiler Tests.....	8vo,	*3 00
— Engine Tests.....	8vo,	*4 00
The above two purchased together.....		*6 00
Barwise, S. The Purification of Sewage.....	12mo,	3 50
Baterden, J. R. Timber. (Westminster Series).....	8vo,	*2 00
Bates, E. L., and Charlesworth, F. Practical Mathematics and Geometry for Technical Students.....	12mo,	*1 50

D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG 5

Beadle, C. Chapters on Papermaking. Five Volumes		
	12mo, each,	*2 00
Beaumont, R. Color in Woven Design.....	8vo,	7 50
— Finishing of Textile Fabrics.....	8vo,	*4 00
Beaumont, W. W. The Steam-Engine Indicator.....	8vo,	2 50
Bedell, F., and Pierce, C. A. Direct and Alternating Current Testing.....	8vo,	*2 00
Beech, F. Dyeing of Cotton Fabrics.....	8vo,	*3 00
— Dyeing of Woolen Fabrics.....	8vo,	*3 50
Beckwith, A. Pottery.....	8vo, paper,	0 60
Begtrup, J. The Slide Valve.....	8vo,	*2 00
Bender, C. E. Continuous Bridges. (Science Series No. 26.)	16mo,	0 50
— Proportions of Piers used in Bridges. (Science Series No. 4.)	16mo,	0 50
Bennett, H.G. The Manufacture of Leather.....	8vo,	*4 50
Bernthsen, A. A Text-book of Organic Chemistry. Trans. by G. M'Gowan.....	12mo,	*2 50
Berry, W. J. Differential Equations of the First Species.	12mo (<i>In Preparation.</i>)	
Bersch, J. Manufacture of Mineral and Lake Pigments. Trans.		
by A. C. Wright.....	8vo,	*5 00
Bertin, L. E. Marine Boilers. Trans. by L. S. Robertson.....	8vo,	5 00
Beveridge, J. Papermaker's Pocket Book.....	12mo,	*4 00
Binns, C. F. Ceramic Technology.....	8vo,	*5 00
— Manual of Practical Potting.....	8vo,	*7 50
— The Potter's Craft.....	12mo,	*2 00
Birchmore, W. H. How to Use a Gas Analysis.....	12mo,	*1 25
Blaine, R. G. The Calculus and Its Applications.....	12mo,	*1 50
Blake, W. H. Brewers' Vade Mecum.....	8vo,	*4 00
Blake, W. P. Report upon the Precious Metals.....	8vo,	2 00
Bligh, W. G. The Practical Design of Irrigation Works.....	8vo,	*6 00
Blyth, A. W. Foods: Their Composition and Analysis.....	8vo,	7 50
— Poisons: Their Effects and Detection.....	8vo,	7 50
Böckmann, F. Celluloid.....	12mo,	*2 50
Bodmer, G. R. Hydraulic Motors and Turbines.....	12mo,	5 00
Boileau, J. T. Traverse Tables.....	8vo,	5 00
Bonney, G. E. The Electro-platers' Handbook.....	12mo,	1 20
Booth, W. H. Water Softening and Treatment.....	8vo,	*2 50

6 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Booth, W. H. Superheaters and Superheating and their Control.....	8vo,	*1 50
Bottcher, A. Cranes: Their Construction, Mechanical Equipment and Working. Trans. by A. Tolhausen....	4to,	*10 00
Bottler, M. Modern Blaching Agents. Trans. by C. Salter.....	12mo,	*2 50
Bottone, S. R. Magnetos for Automobilists.....	12mo,	*1 00
Boulton, S. B. Preservation of Timber. (Science Series No. 82.).....	16mo,	0 50
Bourgougnon, A. Physical Problems. (Science Series No. 113.).....	16mo,	0 50
Bourry, E. Treatise on Ceramic Industries. Trans. by W. P. Rix.....	8vo (In Press.)	
Bow, R. H. A Treatise on Bracing.....	8vo,	1 50
Bowie, A. J., Jr. A Practical Treatise on Hydraulic Mining.....	8vo,	5 00
Bowker, W. R. Dynamo Motor and Switchboard Circuits.....	8vo,	*2 50
Bowles, O. Tables of Common Rocks. (Science Series).....	16mo,	0 50
Bowser, E. A. Elementary Treatise on Analytic Geometry.....	12mo,	1 75
— Elementary Treatise on the Differential and Integral Calculus.....	12mo,	2 25
— Elementary Treatise on Analytic Mechanics.....	12mo,	3 00
— Elementary Treatise on Hydro-mechanics.....	12mo,	2 50
— A Treatise on Roofs and Bridges.....	12mo,	*2 25
Boycott, G. W. M. Compressed Air Work and Diving.....	8vo,	*4 00
Bragg, E. M. Marine Engine Design.....	12mo,	*2 00
Brainard, F. R. The Sextant. (Science Series No. 101.).....	16mo,	
Brassey's Naval Annual for 1910.....	8vo,	*6 00
Brew, W. Three-Phase Transmission.....	8vo,	*2 00
Brewer, R. W. A. The Motor Car.....	12mo,	*2 00
Briggs, R., and Wolff, A. R. Steam-Heating. (Science Series No. 67.).....	16mo,	0 50
Bright, C. The Life Story of Sir Charles Tilson Bright.....	8vo,	*4 50
British Standard Sections.....	8x15	*1 00
Complete list of this series (45 parts) sent on application.		
Broughton, H. H. Electric Cranes and Hoists.....	(In Press.)	
Brown, G. Healthy Foundations. (Science Series No. 80.).....	16mo,	0 50
Brown, H. Irrigation.....	8vo,	*5 00
Brown, Wm. N. The Art of Enamelling on Metal.....	12mo,	*1 00

Brown, Wm. N. Handbook on Japanning and Enamelling.	12mo,	*1 50
— House Decorating and Painting.	12mo,	*1 50
— History of Decorative Art.	12mo,	*1 25
— Dipping, Burnishing, Lacquering and Bronzing Brass Ware.	12mo,	*1 00
— Workshop Wrinkles.	8vo,	*1 00
Browne, R. E. Water Meters. (Science Series No. 81.)	16mo,	0 50
Bruce, E. M. Pure Food Tests.	12mo,	*1 25
Bruhns, Dr. New Manual of Logarithms.	8vo, half mor.,	2 50
Brunner, R. Manufacture of Lubricants, Shoe Polishes and Leather Dressings. Trans. by C. Salter.	8vo,	*3 00
Buel, R. H. Safety Valves. (Science Series No. 21.)	61mo,	0 50
Bulmann, H. F., and Redmayne, R. S. A. Colliery Working and Management.	8vo,	6 00
Burgh, N. P. Modern Marine Engineering.	4to, half mor.,	10 00
Burt, W. A. Key to the Solar Compass.	16mo, leather,	2 50
Burton, F. G. Engineering Estimates and Cost Accounts.	12mo,	*1 50
Buskett, E. W. Fire Assaying.	12mo,	*1 25
Cain, W. Brief Course in the Calculus.	12mo,	*1 75
— Elastic Arches. (Science Series No. 48.)	16mo,	0 50
— Maximum Stresses. (Science Series No. 38.)	16mo,	0 50
— Practical Designing Retaining of Walls. (Science Series No. 3.)	16mo,	0 50
— Theory of Steel-concrete Arches and of Vaulted Structures. (Science Series)	16mo,	0 50
— Theory of Voussoir Arches. (Science Series No. 12.)	16mo,	0 50
— Symbolic Algebra. (Science Series No. 73.)	16mo,	0 50
Campin, F. The Construction of Iron Roofs.	8vo,	2 00
Carpenter, F. D. Geographical Surveying. (Science Series No. 37.)	16mo,	
Carpenter, R. C., and Diederichs, H. Internal-Combustion Engines.	8vo,	*5 00
Carter, E. T. Motive Power and Gearing for Electrical Machinery.	8vo,	*5 00
Carter, H. A. Ramie (Rhea), China Grass.	12mo,	*2 00
Carter, H. R. Modern Flax, Hemp, and Jute Spinning.	8vo,	*3 00
Cathcart, W. L. Machine Design. Part I. Fastenings.	8vo,	*3 00

8 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Cathcart, W. L., and Chaffee, J. I. Elements of Graphic Statics.	8vo,	*3 00
Caven, R. M., and Lander, G. D. Systematic Inorganic Chemistry.....	12mo,	*2 00
Chambers' Mathematical Tables.....	8vo,	1 75
Charnock, G. F. Workshop Practice. (Westminster Series.)	8vo (<i>In Press.</i>)	
Charpentier, P. Timber.....	8vo,	*6 00
Chatley, H. Principles and Designs of Aeroplanes. (Science Series.).....	16mo,	0 50
— How to Use Water Power.....	12mo,	*1 00
Child, C. T. The How and Why of Electricity.....	12mo,	1 00
Christie, W. W. Boiler-waters, Scale, Corrosion, Foaming.....	8vo,	*3 00
— Chimney Design and Theory.....	8vo,	*3 00
— Furnace Draft. (Science Series).....	16mo,	0 50
Church's Laboratory Guide. Rewritten by Edward Kinch..	8vo,	*2 50
Clapperton, G. Practical Papermaking.....	8vo,	2 50
Clark, C. H. Marine Gas Engines.....	(<i>In Press.</i>)	
Clark, D. K. Rules, Tables and Data for Mechanical Engineers	8vo,	5 00
— Fuel: Its Combustion and Economy.....	12mo,	1 50
— The Mechanical Engineer's Pocketbook.....	16mo,	2 00
— Tramways: Their Construction and Working.....	8vo,	7 50
Clark, J. M. New System of Laying Out Railway Turnouts..	12mo,	1 00
Clausen-Thue, W. A B C Telegraphic Code. Fourth Edition	12mo,	*5 00
Fifth Edition.....	8vo,	*7 00
— The A Z Telegraphic Code.....	8vo,	*7 50
Cleemann, T. M. The Railroad Engineer's Practice.....	12mo,	*1 50
Clerk, D., and Idell, F. E. Theory of the Gas Engine. (Science Series No. 62.).....	16mo,	0 50
Clevenger, S. R. Treatise on the Method of Government Surveying.....	16mo, mor.,	2 50
Clouth, F. Rubber, Gutta-Percha, and Balata.....	8vo,	*5 00
Coffin, J. H. C. Navigation and Nautical Astronomy.....	12mo,	*3 50
Colburn, Z., and Thurston, R. H. Steam Boiler Explosions. (Science Series No. 2.).....	16mo,	0 50
Cole, R. S. Treatise on Photographic Optics.....	12mo,	1 50

Coles-Finch, W. Water, Its Origin and Use.....	8vo,	*5 00
Collins, J. E. Useful Alloys and Memoranda for Goldsmiths, Jewelers.....	16mo,	0 50
Constantine, E. Marine Engineers, Their Qualifications and Duties.....	8vo,	*2 00
Coombs, H. A. Gear Teeth. (Science Series No. 120)....	16mo,	0 50
Cooper, W. R. Primary Batteries.....	8vo,	*4 00
—— "The Electrician" Primers.....	8vo,	*5 00
Copperthwaite, W. C. Tunnel Shields.....	4to,	*9 00
Corey, H. T. Water Supply Engineering.....	8vo (<i>In Press.</i>)	
Corfield, W. H. Dwelling Houses. (Science Series No. 50.)	16mo,	0 50
—— Water and Water-Supply. (Science Series No. 17.)....	16mo,	0 50
Cornwall, H. B. Manual of Blow-pipe Analysis.....	8vo,	*2 50
Courtney, C. F. Masonry Dams.....	8vo,	3 50
Cowell, W. B. Pure Air, Ozone, and Water.....	12mo,	*2 00
Craig, T. Motion of a Solid in a Fuel. (Science Series No. 49.)	16mo,	0 50
—— Wave and Vortex Motion. (Science Series No. 43.)....	16mo,	0 50
Cramp, W. Continuous Current Machine Design.....	8vo,	*2 50
Crocker, F. B. Electric Lighting. Two Volumes. 8vo. Vol. I. The Generating Plant.....		3 00
Vol. II. Distributing Systems and Lamps.....		3 00
Crocker, F. B., and Arendt, M. Electric Motors.....	8vo,	*2 50
Crocker, F. B., and Wheeler, S. S. The Management of Electrical Machinery.....	12mo,	*1 00
Cross, C. F., Bevan, E. J., and Sindall, R. W. Wood Pulp and Its Applications. (Westminster Series.).....	8vo (<i>In Press.</i>)	
Crosskey, L. R. Elementary Prospective.....	8vo,	1 00
Crosskey, L. R., and Thaw, J. Advanced Prospective.....	8vo,	1 50
Culley, J. L. Theory of Arches. (Science Series No. 87.)....	16mo,	0 50
Davenport, C. The Book. (Westminster Series.).....	8vo,	*2 00
Davies, D. C. Metalliferous Minerals and Mining.....	8vo,	5 00
—— Earthy Minerals and Mining.....	8vo,	5 00
Davies, E. H. Machinery for Metalliferous Mines.....	8vo,	8 00
Davies, F. H. Electric Power and Traction.....	8vo,	*2 00
Dawson, P. Electric Traction on Railways.....	8vo,	*9 00
Day, C. The Indicator and Its Diagrams.....	12mo,	*2 00

10 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Deerr, N. Sugar and the Sugar Cane.....	8vo,	*3 00
Deite, C. Manual of Soapmaking. Trans. by S. T. King.....	4to,	*5 00
De la Coux, H. The Industrial Uses of Water. Trans. by A. Morris.....	8vo,	*4 50
Del Mar, W. A. Electric Power Conductors.....	8vo,	*2 00
Denny, G. A. Deep-Level Mines of the Rand.....	4to,	*10 00
— Diamond Drilling for Gold.....		*5 00
De Roos, J. D. C. Linkages. (Science Series No. 47.)...	16mo,	0 50
De Varona, A. Sewer Gases. (Science Series No. 55.)...	16mo,	0 50
Derr, W. L. Block Signal Operation.....	Oblong 12mo,	*1 50
Desaint, A. Three Hundred Shades and How to Mix Them.....	8vo,	*10 00
Dibdin, W. J. Public Lighting by Gas and Electricity.....	8vo,	*8 00
— Purification of Sewage and Water.....	8vo,	6 50
Dietrich, K. Analysis of Resins, Balsams, and Gum Resins.....	8vo,	*3 00
Dinger, Lieut. H. C. Care and Operation of Naval Machinery.....	12mo.	*2 00
Dixon, D. B. Machinist's and Steam Engineer's Practical Calculator.....	16mo, mor.,	1 25
Doble, W. A. Power Plant Construction on the Pacific Coast.		
	(In Press.)	
Dodd, G. Dictionary of Manufactures, Mining, Machinery, and the Industrial Arts.....	12mo,	1 50
Dorr, B. F. The Surveyor's Guide and Pocket Table-book.		
	16mo, mor.,	2 00
Down, P. B. Handy Copper Wire Table.....	16mo,	*1 00
Draper, C. H. Elementary Text-book of Light, Heat and Sound.....	12mo,	1 00
— Heat and the Principles of Thermo-dynamics.....	12mo,	1 50
Duckwall, E. W. Canning and Preserving of Food Products.....	8vo,	*5 00
Dumesny, P., and Noyer, J. Wood Products, Distillates, and Extracts.....	8vo,	*4 50
Duncan, W. G., and Penman, D. The Electrical Equipment of Collieries.....	8vo,	*4 00
Duthie, A. L. Decorative Glass Processes. (Westminster Series).....	8vo,	*2 00
Dyson, S. S. Practical Testing of Raw Materials.....	8vo,	*5 00
Eccles, R. G., and Duckwall, E. W. Food Preservatives.....	8vo,	1 00
	paper,	0 50

D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG 11

Eddy, H. T. Researches in Graphical Statics.....	8vo,	1 50
— Maximum Stresses under Concentrated Loads.....	8vo,	1 50
Edgcumbe, K. Industrial Electrical Measuring Instruments.....	8vo,	*2 50
Eissler, M. The Metallurgy of Gold.....	8vo,	7 50
— The Hydrometallurgy of Copper.....	8vo,	*4 50
— The Metallurgy of Silver.....	8vo,	4 00
— The Metallurgy of Argentiferous Lead.....	8vo,	5 00
— Cyanide Process for the Extraction of Gold.....	8vo,	3 00
— A Handbook of Modern Explosives.....	8vo,	5 00
Ekin, T. C. Water Pipe and Sewage Discharge Diagrams..	folio,	*3 00
Eliet, C. W., and Storer, F. H. Compendious Manual of Qualitative Chemical Analysis.....	12mo,	*1 25
Elliot, Major G. H. European Light-house Systems.....	8vo,	5 00
Ennis, Wm. D. Linseed Oil and Other Seed Oils	8vo,	*4 00
— Applied Thermodynamics.....	8vo,	*4 50
— Flying Machines To-day.....	12mo (<i>In Preparation.</i>)	
Erfurt, J. Dyeing of Paper Pulp. Trans. by J. Hubner...	8vo,	*7 50
Erskine-Murray, J. A Handbook of Wireless Telegraphy..	8vo,	*3 50
Evans, C. A. Macadamized Roads.....	(<i>In Press.</i>)	
Ewing, A. J. Magnetic Induction in Iron.....	8vo,	*4 00
Fairie, J. Notes on Lead Ores.....	12mo,	*1 00
— Notes on Pottery Clays.....	12mo,	*1 50
Fairley, W., and Andre, Geo. J. Ventilation of Coal Mines. (Science Series No. 58.).....	16mo,	0 50
Fairweather, W. C. Foreign and Colonial Patent Laws ...	8vo,	*3 00
Fanning, T. T. Hydraulic and Water-supply Engineering.....	8vo,	*5 00
Fauth, P. The Moon in Modern Astronomy. Trans. by J. McCabe.....	8vo,	*2 00
Fay, I. W. The Coal-tar Colors	8vo (<i>In Press.</i>)	
Fernbach, R. L. Glue and Gelatine.....	8vo,	*3 00
Fischer, E. The Preparation of Organic Compounds. Trans. by R. V. Stanford	12mo,	*1 25
Fish, J. C. L. Lettering of Working Drawings.....	Oblong 8vo,	1 00
Fisher, H. K. C., and Darby, W. C. Submarine Cable Testing.....	8vo,	*3 50
Fiske, Lieut. B. A. Electricity in Theory and Practice	8vo,	2 50

12 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Fleischmann, W. The Book of the Dairy. Trans. by C. M. Aikman.....	8vo,	4 00
Fleming, J. A. The Alternate-current Transformer. Two Volumes.....	8vo,	
Vol. I. The Induction of Electric Currents.....		*5 00
Vol. II. The Utilization of Induced Currents.....		*5 00
— Centenary of the Electrical Current.....	8vo,	*0 50
— Electric Lamps and Electric Lighting.....	8vo,	*3 00
— Electric Laboratory Notes and Forms.....	4to,	*5 00
— A Handbook for the Electrical Laboratory and Testing Room. Two Volumes.....	8vo, each,	*5 00
Fluery, H. The Calculus Without Limits or Infinitesimals. Trans. by C. O. Mailloux.....	(In Press.)	
Flynn, P. J. Flow of Water. (Science Series No. 84.)...	16mo,	0 50
— Hydraulic Tables. (Science Series No. 66.).....	16mo,	0 50
Foley, N. British and American Customary and Metric Measures.....	folio,	*3 00
Foster, H. A. Electrical Engineers' Pocket-book. (Sixth Edition.).....	12mo, leather,	5 00
Foster, Gen. J. G. Submarine Blasting in Boston (Mass.) Harbor.....	4to,	3 50
Fowle, F. F. Overhead Transmission Line Crossings....	12mo,	*1 50
— The Solution of Alternating Current Problems.....	8vo (In Press.)	
Fox, W. G. Transition Curves. (Science Series No. 110.)...	16mo,	0 50
Fox, W., and Thomas, C. W. Practical Course in Mechanical Drawing.....	12mo,	1 25
Foye, J. C. Chemical Problems. (Science Series No. 69.)...	16mo,	0 50
— Handbook of Mineralogy. (Science Series No. 86.)...	16mo,	0 50
Francis, J. B. Lowell Hydraulic Experiments.....	4to,	15 00
Frye, A. I. Civil Engineers' Pocket-book.....	(In Press.)	
Fuller, G. W. Investigations into the Purification of the Ohio River.....	4to,	*10 00
Furnell, J. Paints, Colors, Oils, and Varnishes.....	8vo,	*1 00
Gant, L. W. Elements of Electric Traction.....	8vo,	*2 50
Garcke, E., and Fells, J. M. Factory Accounts.....	8vo,	3 00
Garforth, W. E. Rules for Recovering Coal Mines after Explosions and Fires.....	12mo, leather,	1 50

D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG 13

Gaudard, J. Foundations. (Science Series No. 34.).....	16mo,	0 50
Gear, H. B., and Williams, P. F. Electric Central Station Distributing Systems.....	8vo (<i>In Preparation.</i>)	
Geerlign, H. C. P. Cane Sugar and Its Manufacture.....	8vo,	*5 00
Geikie, J. Structural and Field Geology.....	8vo,	*4 00
Gerber, N. Analysis of Milk, Condensed Milk, and Infants' Milk-Food.....	8vo,	1 25
Gerhard, W. P. Sanitation, Water-supply and Sewage Disposal of Country Houses.....	12mo,	*2 00
— Gas Lighting. (Science Series No. 111.).....	16mo,	0 50
— Household Wastes. (Science Series No. 97.).....	16mo,	0 50
— House Drainage. (Science Series No. 63.).....	16mo,	0 50
— Sanitary Drainage of Buildings. (Science Series No. 93.).....	16mo,	0 50
Gerhardi, C. W. H. Electricity Meters.....	8vo,	*4 00
Geschwind, L. Manufacture of Alum and Sulphates. Trans. by C. Salter.....	8vo,	*5 00
Gibbs, W. E. Lighting by Acetylene.....	12mo,	*1 50
— Physics of Solids and Fluids. (Carnegie Technical Schools Text-books.).....		*1 50
Gibson, A. H. Hydraulics and Its Application.....	8vo,	*5 00
— Water Hammer in Hydraulic Pipe Lines.....	12mo,	*2 00
Gilbreth, F. B. Motion Study. A Method for Increasing the Efficiency of the Workman.....	12mo,	*2 00
Gillmore, Gen. Q. A. Limes, Hydraulic Cements and Mortars.....	8vo,	4 00
— Roads, Streets, and Pavements.....	12mo,	2 00
Golding, H. A. The Theta-Phi Diagram.....	12mo,	*1 25
Goldschmidt, R. Alternating Current Commutator Motor.....	8vo,	*3 00
Goodchild, W. Precious Stones. (Westminster Series.).....	8vo,	*2 00
Goodeve, T. M. Textbook on the Steam-engine.....	12mo,	2 00
Gore, G. Electrolytic Separation of Metals.....	8vo,	*3 50
Gould, E. S. Arithmetic of the Steam-engine.....	12mo,	1 00
— Calculus. (Science Series No. 112.).....	16mo,	0 50
— High Masonry Dams. (Science Series No. 22.).....	16mo,	0 50
— Practical Hydrostatics and Hydrostatic Formulas. (Science Series.).....	16mo,	0 50
Grant, J. Brewing and Distilling. (Westminster Series.).....	8vo (<i>In Press.</i>)	
Gray, J. Electrical Influence Machines.....	12mo,	2 00

14 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Greenwood, E. Classified Guide to Technical and Commercial Books.....	8vo,	*3 00
Gregorius, R. Mineral Waxes. Trans. by C. Salter.....	12mo,	*3 00
Griffiths, A. B. A Treatise on Manures.....	12mo,	3 00
— Dental Metallurgy.....	8vo,	*3 50
Gross, E. Hops.....	8vo,	*4 50
Grossman, J. Ammonia and its Compounds.....	12mo,	*1 25
Groth, L. A. Welding and Cutting Metals by Gases or Electricity.....	8vo,	*3 00
Grover, F. Modern Gas and Oil Engines.....	8vo,	*2 00
Gruner, A. Power-loom Weaving.....	8vo,	*3 00
Güldner, Hugo. Internal-Combustion Engines. Trans. by H. Diedrichs.....	4to,	*10 00
Gunther, C. O. Integration.....	12mo,	*1 25
Gurden, R. L. Traverse Tables.....	folio, half mor.	7 50
Guy, A. E. Experiments on the Flexure of Beams.....	8vo,	*1 25
Haeder, H. Handbook on the Steam-engine. Trans. by H. H. P. Powles.....	12mo,	3 00
Hainbach, R. Pottery Decoration. Trans. by C. Slater.....	12mo,	*3 00
Hale, W. J. Calculations of General Chemistry.....	12mo,	*1 00
Hall, C. H. Chemistry of Paints and Paint Vehicles.....	12mo,	*2 00
Hall, R. H. Governors and Governing Mechanism.....	12mo,	*2 00
Hall, W. S. Elements of the Differential and Integral Calculus.....	8vo,	*2 25
— Descriptive Geometry.....	8vo volume and 4to atlas,	*3 50
Haller, G. F., and Cunningham, E. T. The Tesla Coil.....	12mo,	*1 25
Halsey, F. A. Slide Valve Gears.....	12mo,	1 50
— The Use of the Slide Rule. (Science Series.).....	16mo,	0 50
— Worm and Spiral Gearing. (Science Series.).....	16mo,	0 50
Hamilton, W. G. Useful Information for Railway Men.....	16mo,	1 00
Hammer, W. J. Radium and Other Radioactive Substances.....	8vo,	*1 00
Hancock, H. Textbook of Mechanics and Hydrostatics.....	8vo,	1 50
Hardy, E. Elementary Principles of Graphic Statics.....	12mo,	*1 50
Harper, W. B. Utilization of Wood Waste by Distillation.....	4to,	*3 00
Harrison, W. B. The Mechanics' Tool-book.....	12mo,	1 50
Hart, J. W. External Plumbing Work.....	8vo,	*3 00
— Hints to Plumbers on Joint Wiping.....	8vo,	*3 00
— Principles of Hot Water Supply.....	8vo,	*3 00

D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG 15

Hart, J. W. Sanitary Plumbing and Drainage.....	8vo,	*3 00
Haskins, C. H. The Galvanometer and Its Uses.....	16mo,	1 50
Hatt, J. A. H. The Colorist.....	square 12mo,	*1 50
Hausbrand, E. Drying by Means of Air and Steam. Trans. by A. C. Wright.....	12mo,	*2 00
— Evaporating, Condensing and Cooling Apparatus. Trans. by A. C. Wright.....	8vo,	*5 00
Hausner, A. Manufacture of Preserved Foods and Sweetmeats. Trans. by A. Morris and H. Robson.....	8vo,	*3 00
Hawke, W. H. Premier Cipher Telegraphic Code.....	4to,	*5 00
— 100,000 Words Supplement to the Premier Code.....	4to,	*5 00
Hawkesworth, J. Graphical Handbook for Reinforced Concrete Design.....	4to,	*2 50
Hay, A. Alternating Currents.....	8vo,	*2 50
— Principles of Alternate-current Working.....	12mo,	2 00
— Electrical Distributing Networks and Distributing Lines.....	8vo,	*3 50
— Continuous Current Engineering.....	8vo,	*2 50
Heap, Major D. P. Electrical Appliances.....	8vo,	2 00
Heaviside, O. Electromagnetic Theory. Two volumes. 8vo, each,		*5 00
Heck, R. C. H. Steam-Engine and Other Steam Motors. Two Volumes.		
Vol. I. Thermodynamics and the Mechanics.....	8vo,	*3 50
Vol. II. Form, Construction and Working.....	8vo,	*5 00
— Abridged edition of above volumes (Elementary)		
	8vo (In Preparation)	
— Notes on Elementary Kinematics.....	8vo, boards,	*1 00
— Graphics of Machine Forces.....	8vo, boards,	*1 00
Hedges, K. Modern Lightning Conductors.....	8vo,	3 00
Heermann, P. Dyers' Materials. Trans. by A. C. Wright. 12mo,		*2 50
Hellot, Macquer and D'Apligny. Art of Dyeing Wool, Silk and Cotton.....	8vo,	*2 00
Henrici, O. Skeleton Structures.....	8vo,	1 50
Hermann, F. Painting on Glass and Porcelain.....	8vo,	*3 50
Hermann, G. The Graphical Statics of Mechanism. Trans. by A. P. Smith.....	12mo,	2 00
Herzfeld, J. Testing of Yarns and Textile Fabrics.....	8vo,	*3 50
Hildebrandt, A. Airships, Past and Present.....	8vo,	*3 50

16 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Hildenbrand, B. W. Cable-Making. (Science Series No. 32.)	16mo,	0 50
Hildrich, H. Short History of Chemistry.....	(In Press.)	
Hill, J. W. The Purification of Public Water Supplies. New Edition.....	(In Press.)	
— Interpretation of Water Analysis.....	(In Press.)	
Hiroi, I. Plate Girder Construction. (Science Series No. 95.)	16mo,	0 50
— Statically-Indeterminate Stresses.....	12mo,	*2 00
Hirshfeld, C. F. Engineering Thermodynamics. (Science Series.).....	16mo,	0 50
Hobart, H. M. Heavy Electrical Engineering.....	8vo,	*4 50
— Electricity.....	8vo,	*2 00
— Electric Trains.....	8vo,	*2 50
Hobbs, W. R. P. The Arithmetic of Electrical Measurements	12mo,	0 50
Hoff, J. N. Paint and Varnish Facts and Formulas.....	12mo,	*1 50
Hoff, Com.W. B. The Avoidance of Collisions at Sea.	16mo, mor.,	0 75
Hole, W. The Distribution of Gas.....	8vo,	*7 50
Holley, A. L. Railway Practice	folio,	12 00
Holmes, A. B. The Electric Light Popularly Explained.	12mo, paper,	0 50
Hopkins, N. M. Experimental Electrochemistry.....	8vo,	*3 00
— Model Engines and Small Boats.....	12mo,	1 25
Hopkinson, J., Shoolbred, J. N., and Day, R. E. Dynamic Electricity. (Science Series No. 71.).....	16mo,	0 50
Horner, J. Engineers' Turning.....	8vo,	*3 50
— Metal Turning.....	12mo,	1 50
— Toothed Gearing.....	12mo,	2 25
Houghton, C. E. The Elements of Mechanics of Materials.	12mo,	*2 00
Houllevigue, L. The Evolution of the Sciences.....	8vo,	*2 00
Howe, G. Mathematics for the Practical Man.....	12mo,	*1 25
Howorth, J. Repairing and Riveting Glass, China and Earthenware.....	8vo, paper,	*0 50
Hubbard, E. The Utilization of Wood-waste.....	8vo,	*2 50
Humber, W. Calculation of Strains in Girders.....	12mo,	2 50
Humphreys, A. C. The Business Features of Engineering Practice.....	8vo,	*1 25
Hurst, G. H. Handbook of the Theory of Color.....	8vo,	*2 50

D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG 17

Hurst, G.H. Dictionary of Chemicals and Raw Products..	8vo,	*3 00
— Lubricating Oils, Fats and Greases.....	8vo,	*3 00
— Soaps.....	8vo,	*5 00
— Textile Soaps and Oils.....	8vo,	*2 50
Hurst, H. E., and Lattey, R. T. Text-book of Physics.....	8vo,	*3 00
Hutchinson, R. W., Jr. Long Distance Electric Power Transmission.....	12mo,	*3 00
Hutchinson, R. W., Jr., and Ihlseng, M. C. Electricity in Mining.....	12mo (In Press.)	
Hutchinson, W. B. Patents and How to Make Money Out of Them.....	12mo,	1 25
Hutton, W. S. Steam-boiler Construction.....	8vo,	6 00
— Practical Engineer's Handbook.....	8vo,	7 00
— The Works' Manager's Handbook.....	8vo,	6 00
Hyde, E. W. Skew Arches. (Science Series No. 15.)....	16mo,	0 50
Induction Coils. (Science Series No. 53.).....	16mo,	0 50
Ingle, H. Manual of Agricultural Chemistry.....	8vo,	*3 00
Innes, C. H. Problems in Machine Design.....	12mo,	*2 00
— Air Compressors and Blowing Engines.....	12mo,	*2 00
— Centrifugal Pumps.....	12mo,	*2 00
— The Fan.....	12mo,	*2 00
Isherwood, B. F. Engineering Precedents for Steam Machinery.....	8vo,	2 50
Ivatts, E. B. Railway Management at Stations.....	8vo,	*2 50
Jacob, A., and Gould, E. S. On the Designing and Construction of Storage Reservoirs. (Science Series No. 6.)..	16mo,	0 50
Jamieson, A. Text Book on Steam and Steam Engines....	8vo,	3 00
— Elementary Manual on Steam and the Steam Engine.	12mo,	1 50
Jannettaz, E. Guide to the Determination of Rocks. Trans. by G. W. Plympton.....	12mo,	1 50
Jehl, F. Manufacture of Carbons.....	8vo,	*4 00
Jennings, A. S. Commercial Paints and Painting. (Westminster Series.).....	8vo (In Press.)	
Jennison, F. H. The Manufacture of Lake Pigments....	8vo,	*3 00
Jepson, G. Cams and the Principles of their Construction...	8vo,	*1 50
— Mechanical Drawing.....	8vo (In Preparation.)	
Jockin, W. Arithmetic of the Gold and Silversmith.....	12mo,	*1 00

18 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Johnson, G. L. Photographic Optics and Color Photography	8vo,	*3 00
Johnson, W. H. The Cultivation and Preparation of Para Rubber.	8vo,	*3 00
Johnson, W. McA. The Metallurgy of Nickel.....	(In Preparation.)	
Johnston, J. F. W., and Cameron, C. Elements of Agricultural Chemistry and Geology.....	12mo,	2 60
Joly, J. Radioactivity and Geology.....	12mo,	*3 00
Jones, H. C. Electrical Nature of Matter and Radioactivity	12mo,	2 00
Jones, M. W. Testing Raw Materials Used in Paint.....	12mo,	*2 00
Jones, L., and Scard, F. I. Manufacture of Cane Sugar.....	8vo,	*5 00
Joynson, F. H. Designing and Construction of Machine Gear- ing.....	8vo,	2 00
Jüptner, H. F. V. Siderology: The Science of Iron.....	8vo,	*5 00
Kansas City Bridge.....	4to,	6 00
Kapp, G. Alternate Current Machinery. (Science Series No. 96.).....	16mo,	0 50
— Dynamos, Motors, Alternators and Rotary Converters. Trans. by H. H. Simmons.....	8vo,	4 00
— Electric Transmission of Energy.....	12mo,	3 50
Keim, A. W. Prevention of Dampness in Buildings.....	8vo,	*2 00
Keller, S. S. Mathematics for Engineering Students.	12mo, half leather,	
— Algebra and Trigonometry, with a Chapter on Vectors....		*1 75
— Special Algebra Edition.....		*1 00
— Plane and Solid Geometry.....		*1 25
— Analytical Geometry and Calculus.....		*2 00
Kelsey, W. R. Continuous-current Dynamos and Motors.	8vo,	*2 50
Kemble, W. T., and Underhill, C. R. The Periodic Law and the Hydrogen Spectrum.....	8vo, paper,	*0 50
Kemp, J. F. Handbook of Rocks.....	8vo,	*1 50
Kendall, E. Twelve Figure Cipher Code.....	4to,	*15 00
Kennedy, A. B. W., and Thurston, R. H. Kinematics of Machinery. (Science Series No. 54.).....	16mo,	0 50
Kennedy, A. B. W., Unwin, W. C., and Idell, F. E. Compressed Air. (Science Series No. 106.).....	16mo,	0 50

Kennedy, R. Modern Engines and Power Generators. Six Volumes.....	4to,	15 00
Single Volumes.....	each,	3 00
— Electrical Installations. Five Volumes.....	4to,	15 00
Single Volumes.....	each,	3 50
— Flying Machines; Practice and Design.....	12mo,	*2 00
Kennelly, A. E. Electro-dynamic Machinery.....	8vo,	1 50
Kent, W. Strength of Materials. (Science Series No. 41.).....	16mo,	0 50
Kershaw, J. B. C. Fuel, Water and Gas Analysis.....	8vo,	*2 50
— Electrometallurgy. (Westminster Series.).....	8vo,	*2 00
— The Electric Furnace in Iron and Steel Production.....	12mo,	*1 50
Kingdon, J. A. Applied Magnetism.....	8vo,	*3 00
Kinzbrunner, C. Alternate Current Windings.....	8vo,	*1 50
— Continuous Current Armatures.....	8vo,	*1 50
— Testing of Alternating Current Machines.....	8vo,	*2 00
Kirkaldy, W. G. David Kirkaldy's System of Mechanical Testing.....	4to,	10 00
Kirkbride, J. Engraving for Illustration.....	8vo,	*1 50
Kirkwood, J. P. Filtration of River Waters.....	4to,	7 50
Klein, J. F. Design of a High speed Steam-engine.....	8vo,	*5 00
— Physical Significance of Entropy.....	8vo,	*1 50
Kleinhans, F. B. Boiler Construction.....	8vo,	3 00
Knight, Capt. A. M. Modern Steamship.....	8vo,	*7 50
	Half Mor.	*9 00
Knox, W. F. Logarithm Tables.....	(In Preparation.)	
Knott, C. G., and Mackay, J. S. Practical Mathematics.....	8vo,	2 00
Koester, F. Steam-Electric Power Plants.....	4to,	*5 00
— Hydroelectric Developments and Engineering.....	4to,	*5 00
Koller, T. The Utilization of Waste Products.....	8vo,	*3 50
— Cosmetics.....	8vo,	*2 50
Krauch, C. Testing of Chemical Reagents. Trans. by J. A. Williamson and L. W. Dupre.....	8vo,	*3 00
Lambert, T. Lead and its Compounds.....	8vo,	*3 50
— Bone Products and Manures.....	8vo,	*3 00
Lamborn, L. L. Cottonseed Products.....	8vo,	*3 00
— Modern Soaps, Candles, and Glycerin.....	8vo,	*7 50
Lamprecht, R. Recovery Work After Pit Fires. Trans. by C. Salter.....	8vo,	*4 00

20 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Lanchester, F. W. Aerial Flight. Two Volumes. 8vo.		
Vol. I. Aerodynamics.....		*6 00
Vol. II. Aerodnetics.....		*6 00
Larner, E. T. Principles of Alternating Currents.....	12mo,	*1 25
Larrabee, C. S. Cipher and Secret Letter and Telegraphic Code.....	16mo,	0 60
La Rue, B. F. Swing Bridges. (Science Series No. 107.)	16mo,	0 50
Lassar-Cohn, Dr. Modern Scientific Chemistry. Trans. by M. M. Pattison Muir.....	12mo,	*2 00
Latimer, L. H., Field, C. J., and Howell, J. W. Incandescent Electric Lighting. (Science Series No. 57.)	16mo,	0 50
Latta, M. N. Handbook of American Gas-Engineering Practice.	8vo,	*4 50
— American Producer Gas Practice	4to,	*6 00
Leask, A. R. Breakdowns at Sea.....	12mo,	2 00
— Triple and Quadruple Expansion Engines.....	12mo,	2 00
— Refrigerating Machinery	12mo,	2 00
Lecky, S. T. S. "Wrinkles" in Practical Navigation	8vo,	*8 00
Le Doux, M. Ice-Making Machines. (Science Series No. 46.)	16mo,	0 50
Leeds, C. C. Mechanical Drawing for Trade Schools. oblong, 4to,		
High School Edition.....		*1 25
Machinery Trades Edition.....		*2 00
Lefèvre, L. Architectural Pottery. Trans. by H. K. Bird and W. M. Binns.....	4to,	*7 50
Lehner, S. Ink Manufacture. Trans. by A. Morris and H. Robson.....	8vo,	*2 50
Lemstrom, S. Electricity in Agriculture and Horticulture.	8vo,	*1 50
Le Van, W. B. Steam-Engine Indicator. (Science Series No. 78.).....	16mo,	0 50
Lewes, V. B. Liquid and Gaseous Fuels. (Westminster Series.)	8vo,	*2 00
Lieber, B. F. Lieber's Standard Telegraphic Code	8vo,	*10 00
— Code. German Edition.....	8vo,	*10 00
— Spanish Edition.....	8vo,	*10 00
— French Edition.....	8vo,	*10 00
— Terminal Index.....	8vo,	*2 50
— Lieber's Appendix.....	folio,	*15 00

D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG 21

Lieber, B. F. Handy Tables.....	4to,	*2 50
— Bankers and Stockbrokers' Code and Merchants and Shippers' Blank Tables.....	8vo,	*15 00
— 100,000,000 Combination Code.....	8vo,	*15 00
— Engineering Code.....	8vo,	*10 00
Livermore, V. P., and Williams, J. How to Become a Competent Motorman.....	12mo,	*1 00
Livingstone, R. Design and Construction of Commutators.....	8vo,	*2 25
Lobben, P. Machinists' and Draftsmen's Handbook.....	8vo,	2 50
Locke, A. G. and C. G. Manufacture of Sulphuric Acid.....	8vo,	10 00
Lockwood, T. D. Electricity, Magnetism, and Electro-teleg-raphy.....	8vo,	2 50
— Electrical Measurement and the Galvanometer.....	12mo,	1 50
Lodge, O. J. Elementary Mechanics.....	12mo,	1 50
— Signalling Across Space without Wires.....	8vo,	*2 00
Lord, R. T. Decorative and Fancy Fabrics.....	8vo,	*3 50
Loring, A. E. A Handbook of the Electromagnetic Telegraph. (Science Series No. 39).....	16mo,	0 50
Loewenstein, L. C., and Crissey, C. P. Centrifugal Pumps. (<i>In Press.</i>)		
Lucke, C. E. Gas Engine Design.....	8vo,	*3 00
— Power Plants: their Design, Efficiency, and Power Costs. 2 vols.....		(<i>In Preparation.</i>)
— Power Plant Papers. Form I. The Steam Power Plant	paper,	*1 50
Lunge, G. Coal-tar Ammonia. Two Volumes.....	8vo,	*15 00
— Manufacture of Sulphuric Acid and Alkali. Three Volumes	8vo,	
Vol. I. Sulphuric Acid. In two parts.....		*15 00
Vol. II. Salt Cake, Hydrochloric Acid and Leblanc Soda. In two parts.....		*15 00
Vol. III. Ammonia Soda.....		*15 00
— Technical Chemists' Handbook.....	12mo, leather,	*3 50
— Technical Methods of Chemical Analysis. Trans. by C. A. Keane. In collaboration with the corps of specialists.		
Vol. I. In two parts.....	8vo,	*15 00
Vols. II and III.....		(<i>In Preparation.</i>)
Lupton, A., Parr, G. D. A., and Perkin, H. Electricity as Applied to Mining.....	8vo,	*4 50
Luquer, L. M. Minerals in Rock Sections.....	8vo,	*1 50

22 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Macewen, H. A. Food Inspection.....	8vo,	*2 50
Mackenzie, N. F. Notes on Irrigation Works.....	8vo,	*2 50
Mackie, J. How to Make a Woolen Mill Pay.....	8vo,	*2 00
Mackrow, C. Naval Architect's and Shipbuilder's Pocket-book.....	16mo, leather,	5 00
Maguire, Capt. E. The Attack and Defense of Coast Fortifications.....	8vo,	2 50
Maguire, Wm. R. Domestic Sanitary Drainage and Plumbing.....	8vo,	4 00
Mallet, A. Compound Engines. Trans. by R. R. Buel. (Science Series No. 10.).....	16mo,	
Mansfield, A. N. Electro-magnets. (Science Series No. 64).....	16mo,	0 50
Marks, E. C. R. Construction of Cranes and Lifting Machinery.....	12mo,	*1 50
— Construction and Working of Pumps.....	12mo,	*1 50
— Manufacture of Iron and Steel Tubes.....	12mo,	*2 00
— Mechanical Engineering Materials.....	12mo,	*1 00
Marks, G. C. Hydraulic Power Engineering.....	8vo,	3 50
— Inventions, Patents and Designs.....	12mo,	*1 00
Markham, E. R. The American Steel Worker.....	12mo,	2 50
Marlow, T. G. Drying Machinery and Practice.....	8vo,	*5 00
Marsh, C. F. Concise Treatise on Reinforced Concrete....	8vo,	*2 50
Marsh, C. F., and Dunn, W. Reinforced Concrete.....	4to,	*5 00
— Manual of Reinforced Concrete and Concrete Block Construction.....	16mo, mor.,	*2 50
Marshall, W. J., and Sankey, H. R. Gas Engines. (Westminster Series.).....	8vo,	*2 00
Massie, W. W., and Underhill, C. R. Wireless Telegraphy and Telephony.....	12mo,	*1 00
Matheson, D. Australian Saw-Miller's Log and Timber Ready Reckoner.....	12mo, leather,	1 50
Mathot, R. E. Internal Combustion Engines.....	8vo,	*6 00
Maurice, W. Electric Blasting Apparatus and Explosives ..	8vo,	*3 50
— Shot Firer's Guide.....	8vo,	*1 50
Maxwell, J. C. Matter and Motion. (Science Series No. 36.).....	16mo,	0 50
Maxwell, W. H., and Brown, J. T. Encyclopedia of Municipal and Sanitary Engineering.....	4to,	*10 00
Mayer, A. M. Lecture Notes on Physics.....	8vo,	2 00

D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG 23

McCullough, R. S. Mechanical Theory of Heat.....	8vo,	3 50
McIntosh, J. G. Technology of Sugar.....	8vo,	*4 50
— Industrial Alcohol.....	8vo,	*3 00
— Manufacture of Varnishes and Kindred Industries. Three Volumes. 8vo.		
Vol. I. Oil Crushing, Refining and Boiling		*3 50
Vol. II. Varnish Materials and Oil Varnish Making.....		*4 00
Vol. III.....	(In Preparation.)	
McKnight, J. D., and Brown, A. W. Marine Multitubular Boilers.....		*1 50
McMaster, J. B. Bridge and Tunnel Centres. (Science Series No. 20.).....	16mo,	0 50
McMechen, F. L. Tests for Ores, Minerals and Metals....	12mo,	*1 00
McNeill, B. McNeill's Code.....	8vo,	*6 00
McPherson, J. A. Water-works Distribution	8vo,	2 50
Melick, C. W. Dairy Laboratory Guide.....	12mo,	*1 25
Merck, E. Chemical Reagents; Their Purity and Tests....	8vo,	*1 50
Merritt, Wm. H. Field Testing for Gold and Silver. 16mo, leather,		1 50
Meyer, J. G. A., and Pecker, C. G. Mechanical Drawing and Machine Design.....	4to,	5 00
Michell, S. Mine Drainage.....	8vo,	10 00
Mierzinski, S. Waterproofing of Fabrics. Trans. by A. Morris and H. Robson.....	8vo,	*2 50
Miller, E. H. Quantitative Analysis for Mining Engineers. 8vo,		*1 50
Miller, G. A. Determinants. (Science Series No. 105.)..	16mo,	
Milroy, M. E. W. Home Lace-making.....	12mo,	*1 00
Minifie, W. Mechanical Drawing.....	8vo,	*4 00
Mitchell, C. A., and Prideaux, R. M. Fibres Used in Textile and Allied Industries.....	8vo,	
Modern Meteorology	12mo,	1 50
Monckton, C. C. F. Radiotelegraphy. (Westminster Series.) 8vo,		*2 00
Monteverde, R. D. Vest Pocket Glossary of English-Spanish, Spanish-English Technical Terms.....	64mo, leather,	*1 00
Moore, E. C. S. New Tables for the Complete Solution of Ganguillet and Kutter's Formula	8vo,	*5 00
Moreing, C. A., and Neal, T. New General and Mining Tele- graph Code.....	8vo,	*5 00
Morgan, A. P. Wireless Telegraph Construction for Amateurs. 12mo,		*1 50

24 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Moses, A. J. The Characters of Crystals	8vo,	*2 00
Moses, A. J., and Parsons, C. I. Elements of Mineralogy	8vo,	*2 50
Moss, S. A. Elements of Gas Engine Design. (Science Series.)	16mo,	0 50
— The Lay-out of Corliss Valve Gears. (Science Series.)	16mo,	0 50
Mullin, J. P. Modern Moulding and Pattern-making	12mo,	2 50
Munby, A. E. Chemistry and Physics of Building Materials. (Westminster Series.)	8vo,	*2 00
Murphy, J. G. Practical Mining	16mo,	1 00
Murray, J. A. Soils and Manures. (Westminster Series.)	8vo,	*2 00
Naquet, A. Legal Chemistry	12mo,	2 00
Nasmith, J. The Student's Cotton Spinning	8vo,	3 00
Neilson, R. M. Aeroplane Patents	8vo,	*2 00
Nerz, F. Searchlights. Trans. by C. Rodgers	8vo,	*3 00
Neuberger, H., and Noalhat, H. Technology of Petroleum. Trans. by J. G. McIntosh	8vo,	*10 00
Newall, J. W. Drawing, Sizing and Cutting Bevel-gears.	8vo,	1 50
Newlands, J. Carpenters and Joiners' Assistant	folio, half mor.,	15 00
Nicol, G. Ship Construction and Calculations	8vo,	*4 50
Nipher, F. E. Theory of Magnetic Measurements	12mo,	1 00
Nisbet, H. Grammar of Textile Design	8vo,	*3 00
Nolan, H. The Telescope. (Science Series No. 51.)	16mo,	0 50
Noll, A. How to Wire Buildings	12mo,	1 50
Nugent, E. Treatise on Optics	12mo,	1 50
O'Connor, H. The Gas Engineer's Pocketbook	12mo, leather,	3 50
— Petrol Air Gas	12mo,	*0 75
Ohm, G. S., and Lockwood, T. D. Galvanic Circuit. Trans. by William Francis. (Science Series No. 102.)	16mo,	0 50
Olsen, J. C. Text book of Quantitative Chemical Analysis	8vo,	*4 00
Olsson, A. Motor Control, in Turret Turning and Gun Elevating. (U. S. Navy Electrical Series, No. 1.)	12mo, paper,	*0 50
Oudin, M. A. Standard Polyphase Apparatus and Systems	8vo,	*3 00
Palaz, A. Industrial Photometry. Trans. by G. W. Patterson, Jr.	8vo,	*4 00

D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG 25

Parnely, C. Colliery Manager's Handbook.....	8vo,	*10 00
Parr, G. D. A. Electrical Engineering Measuring Instruments.	8vo,	*3 50
Parry, E. J. Chemistry of Essential Oils and Artificial Per- fumes.....	8vo,	*5 00
Parry, E. J., and Coste, J. H. Chemistry of Pigments	8vo,	*4 50
Parry, L. A. Risk and Dangers of Various Occupations.....	8vo,	*3 00
Parshall, H. F., and Hobart, H. M. Armature Windings.....	4to,	*7 50
—— Electric Railway Engineering.....	4to,	*10 00
Parshall, H. F., and Parry, E. Electrical Equipment of Tram- ways.....	(In Press.)	
Parsons, S. J. Malleable Cast Iron.....	8vo,	*2 50
Passmore, A. C. Technical Terms Used in Architecture	8vo,	*3 50
Patterson, D. The Color Printing of Carpet Yarns.....	8vo,	*3 50
—— Color Matching on Textiles.....	8vo,	*3 00
—— The Science of Color Mixing.....	8vo,	*3 00
Patton, H. B. Lecture Notes on Crystallography.....	8vo,	*1 25
Paulding, C. P. Condensation of Steam in Covered and Bare Pipes.....	8vo,	*2 00
—— Transmission of Heat through Cold-storage Insulation	12mo,	*1 00
Peirce, B. System of Analytic Mechanics.....	4to,	10 00
Pendred, V. The Railway Locomotive. (Westminster Series.)	8vo,	*2 00
Perkin, F. M. Practical Methods of Inorganic Chemistry.	12mo,	*1 00
Perrigo, O. E. Change Gear Devices.....	8vo,	1 00
Perrine, F. A. C. Conductors for Electrical Distribution	8vo,	*3 50
Perry, J. Applied Mechanics.....	8vo,	*2 50
Petit, G. White Lead and Zinc White Paints.....	8vo,	*1 50
Petit, R. How to Build an Aeroplane. Trans. by T. O'B. Hubbard, and J. H. Ledeboer.....	8vo,	*1 50
Pettit, Lieut. J. S. Graphic Processes. (Science Series No. 76.)	16mo,	0 50
Philbrick, P. H. Beams and Girders. (Science Series No. 88.)	16mo,	
Phillips, J. Engineering Chemistry.....	8vo,	*4 50
—— Gold Assaying.....	8vo,	*2 50
—— Dangerous Goods.....	8vo,	3 50

26 D. VAN NOSTRAND COMPANY'S SHORT TITLE CATALOG

Phin, J. Seven Follies of Science.....	12mo,	*1 25
— Household Pests, and How to Get Rid of Them	8vo (<i>In Preparation.</i>)	
Pickworth, C. N. The Indicator Handbook. Two Volumes	12mo, each,	1 50
— Logarithms for Beginners	12mo, boards,	0 50
— The Slide Rule.....	12mo,	1 00
Plane Table, The	8vo,	2 00
Plattner's Manual of Blowpipe Analysis. Eighth Edition, re- vised. Trans. by H. B. Cornwall.....	8vo,	*4 00
Plympton, G. W. The Aneroid Barometer. (Science Series.)	16mo,	0 50
— How to become an Engineer. (Science Series No. 100.)	16mo,	0 50
— Van Nostrand's Table Book. (Science Series No. 104.)	16mo,	0 50
Pochet, M. L. Steam Injectors. Translated from the French. (Science Series No. 29.).....	16mo,	0 50
Pocket Logarithms to Four Places. (Science Series.).....	16mo,	0 50
	leather,	1 00
Pope, F. L. Modern Practice of the Electric Telegraph....	8vo,	1 50
Popplewell, W. C. Elementary Treatise on Heat and Heat Engines	12mo,	*3 00
— Prevention of Smoke.....	8vo,	*3 50
— Strength of Minerals.....	8vo,	*1 75
Potter, T. Concrete.....	8vo,	*3 00
Practical Compounding of Oils, Tallow and Grease.....	8vo,	*3 50
Practical Iron Founding	12mo,	1 50
Pray, T., Jr. Twenty Years with the Indicator.....	8vo,	2 50
— Steam Tables and Engine Constant.....	8vo,	2 00
— Calorimeter Tables.....	8vo,	1 00
Preece, W. H. Electric Lamps.....	(<i>In Press.</i>)	
Prelini, C. Earth and Rock Excavation.....	8vo,	*3 00
— Dredges and Dredging	8vo (<i>In Press.</i>)	
— Graphical Determination of Earth Slopes.....	8vo,	*2 00
— Tunneling	8vo,	3 00
Prescott, A. B. Organic Analysis.....	8vo,	5 00
Prescott, A. B., and Johnson, O. C. Quauitative Chemical Analysis.....	8vo,	*3 50

Prescott, A. B., and Sullivan, E. C. First Book in Qualitative Chemistry.....	12mo,	*1 50
Pritchard, O. G. The Manufacture of Electric-light Carbons. 8vo, paper,		*0 60
Prost, E. Chemical Analysis of Fuels, Ores, Metals. Trans. by J. C. Smith.....	8vo,	*4 50
Pullen, W. W. F. Application of Graphic Methods to the Design of Structures.....	12mo,	*2 50
— Injectors: Theory, Construction and Working.....	12mo,	*1 50
Pulsifer, W. H. Notes for a History of Lead.....	8vo,	4 00
Purchase, W. R. Masonry.....	12mo,	*3 00
Putsch, A. Gas and Coal-dust Firing.....	8vo,	*3 00
Pynchon, T. R. Introduction to Chemical Physics.....	8vo,	3 00
Rafter, G. W. Mechanics of Ventilation. (Science Series No. 33.).....	16mo,	0 50
— Potable Water. (Science Series No. 103.).....	16mo,	0 50
— Treatment of Septic Sewage. (Science Series.).....	16mo,	0 50
Rafter, G. W., and Baker, M. N. Sewage Disposal in the United States.....	4to,	*6 00
Raikes, H. P. Sewage Disposal Works.....	8vo,	*4 00
Railway Shop Up-to-Date.....	4to,	2 00
Ramp, H. M. Foundry Practice.....	(In Press.)	
Randall, P. M. Quartz Operator's Handbook.....	12mo,	2 00
Randau, P. Enamels and Enamelling.....	8vo,	*4 00
Rankine, W. J. M. Applied Mechanics.....	8vo,	5 00
— Civil Engineering.....	8vo,	6 50
— Machinery and Millwork.....	8vo,	5 00
— The Steam-engine and Other Prime Movers.....	8vo,	5 00
— Useful Rules and Tables.....	8vo,	4 00
Rankine, W. J. M., and Bamber, E. F. A Mechanical Text-book.....	8vo,	3 50
Raphael, F. C. Localization of Faults in Electric Light and Power Mains.....	8vo,	*3 00
Rathbone, R. L. B. Simple Jewellery.....	8vo,	*2 00
Rateau, A. Flow of Steam through Nozzles and Orifices. Trans. by H. B. Brydon.....	8vo,	*1 50
Rausenberger, F. The Theory of the Recoil of Guns.....	8vo,	*4 50

28 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Rautenstrauch, W. Notes on the Elements of Machine Design,	8vo, boards,	*1	50
Rautenstrauch, W., and Williams, J. T. Machine Drafting and Empirical Design.			
Part I. Machine Drafting.....	8vo,	*1	25
Part II. Empirical Design.....	(In Preparation.)		
Raymond, E. B. Alternating Current Engineering.....	12mo,	*2	50
Rayner, H. Silk Throwing and Waste Silk Spinning.....	8vo,	*2	50
Recipes for the Color, Paint, Varnish, Oil, Soap and Drysaltery Trades.....	8vo,	*3	50
Recipes for Flint Glass Making.....	12mo,	*4	50
Redwood, B. Petroleum. (Science Series No. 92.).....	16mo,	0	50
Reed's Engineers' Handbook.....	8vo,	*5	00
— Key to the Nineteenth Edition of Reed's Engineers' Handbook.....	8vo,	*3	00
— Useful Hints to Sea-going Engineers.....	12mo,	1	50
— Marine Boilers.....	12mo,	2	00
Reinhardt, C. W. Lettering for Draftsmen, Engineers, and Students.....	oblong 4to, boards,	1	00
— The Technic of Mechanical Drafting... oblong 4to, boards,		*1	00
Reiser, F. Hardening and Tempering of Steel. Trans. by A. Morris and H. Robson.....	12mo,	*2	50
Reiser, N. Faults in the Manufacture of Woollen Goods. Trans. by A. Morris and H. Robson.....	8vo,	*2	50
— Spinning and Weaving Calculations.....	8vo,	*5	00
Renwick, W. G. Marble and Marble Working.....	8vo,	5	00
Reynolds, O., and Idell, F. E. Triple Expansion Engines. (Science Series No. 99.).....	16mo,	0	50
Rhead, G. F. Simple Structural Woodwork.....	12mo,	*1	00
Rhead, G. W. British Pottery Marks.....	8vo,	*3	00
Rice, J. M., and Johnson, W. W. A New Method of Obtaining the Differential of Functions.....	12mo,	0	50
Richardson, J. The Modern Steam Engine.....	8vo,	*3	50
Richardson, S. S. Magnetism and Electricity.....	12mo,	*2	00
Rideal, S. Glue and Glue Testing.....	8vo,	*4	00
Rings, F. Concrete in Theory and Practice.....	12mo,	*2	50
Ripper, W. Course of Instruction in Machine Drawing... folio,		*6	00
Roberts, F. C. Figure of the Earth. (Science Series No. 79.)	16mo,	0	50

D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG 29

Roberts, J., Jr. Laboratory Work in Electrical Engineering.	8vo,	*2 00
Robertson, L. S. Water-tube Boilers.....	8vo,	3 00
Robinson, J. B. Architectural Composition.....	8vo,	*2 50
Robinson, S. W. Practical Treatise on the Teeth of Wheels. (Science Series No. 24.).....	16mo,	0 50
—— Railroad Economics. (Science Series No. 59.).....	16mo,	0 50
—— Wrought Iron Bridge Members. (Science Series No. 60.)	16mo,	0 50
Robson, J. H. Machine Drawing and Sketching.....	(In Press.)	
Roebling, J. A. Long and Short Span Railway Bridges..	folio,	25 00
Rogers, A. A Laboratory Guide of Industrial Chemistry..	12mo,	*1 50
Rogers, A., and Aubert, A. B. Industrial Chemistry.....	(In Press.)	
Rogers, F. Magnetism of Iron Vessels. (Science Series No. 30.)	16mo,	0 50
Rollins, W. Notes on X-Light.....	8vo,	5 00
Rose, J. The Pattern-makers' Assistant.....	8vo,	2 50
—— Key to Engines and Engine-running.....	12mo,	2 50
Rose, T. K. The Precious Metals. (Westminster Series.)	8vo,	*2 00
Rosenhain, W. Glass Manufacture. (Westminster Series.)	8vo,	*2 00
Ross, W. A. Blowpipe in Chemistry and Metallurgy..	12mo,	*2 00
Rossiter, J. T. Steam Engines. (Westminster Series.)	8vo (In Press.)	
—— Pumps and Pumping Machinery. (Westminster Series.)	8vo (In Press.)	
Roth. Physical Chemistry.....	8vo,	*2 00
Rouillion, L. The Economics of Manual Training.....	8vo,	2 00
Rowan, F. J. Practical Physics of the Modern Steam-boiler	8vo,	7 50
Rowan, F. J., and Idell, F. E. Boiler Incrustation and Corro- sion. (Science Series No. 27.).....	16mo,	0 50
Roxburgh, W. General Foundry Practice.....	8vo,	*3 50
Ruhmer, E. Wireless Telephony. Trans. by J. Erskine- Murray.....	8vo,	*3 50
Russell, A. Theory of Electric Cables and Networks.....	8vo,	*3 00
Sabine, R. History and Progress of the Electric Telegraph.	12mo,	1 25
Saeltzer, A. Treatise on Acoustics.....	12mo,	1 00

30 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Salomons, D. Electric Light Installations. 12mo.		
Vol. I. The Management of Accumulators.....	2	50
Vol. II. Apparatus.....	2	25
Vol. III. Applications.....	1	50
Sanford, P. G. Nitro-explosives.....	8vo,	*4 00
Saunders, C. H. Handbook of Practical Mechanics.....	16mo,	1 00
	leather,	1 25
Saunniar, C. Watchmaker's Handbook.....	12mo,	3 00
Sayers, H. M. Brakes for Tram Cars.....	8vo,	*1 25
Scheele, C. W. Chemical Essays.....	8vo,	*2 00
Schellen, H. Magneto-electric and Dynamo-electric Machines		
	8vo,	5 00
Scherer, R. Casein. Trans. by C. Salter.....	8vo,	*3 00
Schmall, C. N. First Course in Analytic Geometry, Plane and		
Solid.....	12mo, half leather,	*1 75
Schmall, C. N., and Schack, S. M. Elements of Plane Geometry		
	12mo,	*1 25
Schmeer, L. Flow of Water.....	8vo,	*3 00
Schumann, F. A Manual of Heating and Ventilation.		
	12mo, leather,	1 50
Schwartz, E. H. L. Causal Geology.....	8vo,	*2 50
Schweizer, V., Distillation of Resins.....	8vo,	*3 50
Scott, W. W. Qualitative Chemical Analysis. A Laboratory		
Manual.....	8vo	*1 50
Scribner, J. M. Engineers' and Mechanics' Companion.		
	16mo, leather,	1 50
Searle, G. M. "Sumners' Method." Condensed and Improved.		
(Science Series No. 124.).....	8vo,	6 00
Seaton, A. E. Manual of Marine Engineering.....	8vo,	6 00
Seaton, A. E., and Rounthwaite, H. M. Pocket-book of Marine		
Engineering.....	16mo, leather,	3 00
Seeligmann, T., Torrilhon, G. L., and Falconnet, H. India		
Rubber and Gutta Percha. Trans. by J. G. McIntosh		
	8vo,	*5 00
Seidell, A. Solubilities of Inorganic and Organic Substances		
	8vo,	*3 00
Sellew, W. H. Steel Rails.....	4to (<i>In Press.</i>)	
Senter, G. Outlines of Physical Chemistry.....	12mo,	*1 50
Sever, G. F. Electric Engineering Experiments	8vo, boards,	*1 00

Sever, G. F., and Townsend, F. Laboratory and Factory Tests in Electrical Engineering.....	8vo,	*2 50
Sewall, C. H. Wireless Telegraphy.....	8vo,	*2 00
—— Lessons in Telegraphy.....	12mo,	*1 00
Sewell, T. Elements of Electrical Engineering.....	8vo,	*3 00
—— The Construction of Dynamos.....	8vo,	*3 00
Sexton, A. H. Fuel and Refractory Materials.....	12mo,	*2 50
—— Chemistry of the Materials of Engineering.....	12mo,	*2 50
—— Alloys (Non-Ferrous).....	8vo,	*3 00
—— The Metallurgy of Iron and Steel.....	8vo,	*6 50
Seymour, A. Practical Lithography.....	8vo,	*2 50
—— Modern Printing Inks.....	8vo,	*2 00
Shaw, H. S. H. Mechanical Integrators. (Science Series No. 83.).....	16mo,	0 50
Shaw, P. E. Course of Practical Magnetism and Electricity.	8vo,	*1 00
Shaw, S. History of the Staffordshire Potteries.....	8vo,	*3 00
—— Chemistry of Compounds Used in Porcelain Manufacture	8vo,	*5 00
Sheldon, S., and Hausmann, E. Direct Current Machines.	8vo,	*2 50
—— Alternating-current Machines.....	8vo,	*2 50
Sherer, R. Casein. Trans. by C. Salter.....	8vo,	*3 00
Sherriff, F. F. Oil Merchants' Manual.....	12mo,	*3 50
Shields, J. E. Notes on Engineering Construction.....	12mo,	1 50
Shock, W. H. Steam Boilers.....	4to, half mor.,	15 00
Shreve, S. H. Strength of Bridges and Roofs.....	8vo,	3 50
Shunk, W. F. The Field Engineer.....	12mo, mor.,	2 50
Simmons, W. H., and Appleton, H. A. Handbook of Soap Manufacture.....	8vo,	*3 00
Simms, F. W. The Principles and Practice of Leveling.....	8vo,	2 50
—— Practical Tunneling.....	8vo,	7 50
Simpson, G. The Naval Constructor.....	12mo, mor.,	*5 00
Sinclair, A. Development of the Locomotive Engine.	8vo, half leather,	5 00
Sindall, R. W. Manufacture of Paper. (Westminster Series.)	8vo,	*2 00
Sloane, T. O'C. Elementary Electrical Calculations.....	12mo,	*2 00
Smith, C. A. M. Handbook of Testing. Vol. I. Materials. (In Press.)		

32 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Smith, C. F. Practical Alternating Currents and Testing	8vo,	*2 50
— Practical Testing of Dynamos and Motors	8vo,	*2 00
Smith, F. E. Handbook of General Instruction for Mechanics	12mo,	1 50
Smith, I. W. The Theory of Deflections and of Latitudes and Departures	16mo, mor.,	3 00
Smith, J. C. Manufacture of Paint	8vo,	*3 00
Smith, W. Chemistry of Hat Manufacturing	12mo,	*3 00
Snell, A. T. Electric Motive Power	8vo,	*4 00
Snow, W. G. Pocketbook of Steam Heating and Ventilation	<i>(In Press.)</i>	
Snow, W. G., and Nolan, T. Ventilation of Buildings. (Science Series No. 5.)	16mo,	0 50
Soddy, F. Radioactivity	8vo,	*3 00
Solomon, M. Electric Lamps. (Westminster Series.)	8vo,	*2 00
Sothorn, J. W. The Marine Steam Turbine	8vo,	*5 00
Soxhlet, D. H. Dyeing and Staining Marble. Trans. by A. Morris and H. Robson	8vo,	*2 50
Spang, H. W. A Practical Treatise on Lightning Protection	12mo,	1 00
Spangenburg, L. Fatigue of Metals. Translated by S. H. Shreve. (Science Series No. 23.)	16mo,	0 50
Specht, G. J., Hardy, A. S., McMaster, J. B., and Walling. Topographical Surveying. (Science Series No. 72.)	16mo,	0 50
Speyers, C. L. Text-book of Physical Chemistry	8vo,	*2 25
Stahl, A. W. Transmission of Power. (Science Series No. 28.)	16mo,	
Stahl, A. W., and Woods, A. T. Elementary Mechanism	12mo,	*2 00
Staley, C., and Pierson, G. S. The Separate System of Sewerage	8vo,	*3 00
Standage, H. C. Leatherworkers' Manual	8vo,	*3 50
— Sealing Waxes, Wafers, and Other Adhesives	8vo,	*2 00
— Agglutinants of all Kinds for all Purposes	12mo,	*3 50
Stansbie, J. H. Iron and Steel. (Westminster Series.)	8vo,	*2 00
Stevens, H. P. Paper Mill Chemist	16mo,	*2 50
Stevenson, J. L. Blast-Furnace Calculations	12mo, leather,	*2 00
Stewart, A. Modern Polyphase Machinery	12mo,	*2 00
Stewart, G. Modern Steam Traps	12mo,	*1 25
Stiles, A. Tables for Field Engineers	12mo,	1 00

Stillman, P. Steam-engine Indicator.....	12mo,	1 00
Stodola, A. Steam Turbines. Trans. by L. C. Loewenstein.	8vo,	*5 00
Stone, H. The Timbers of Commerce.....	8vo,	3 50
Stone, Gen. R. New Roads and Road Laws.....	12mo,	1 00
Stopes, M. Ancient Plants.....	8vo,	*2 00
— The Study of Plant Life	8vo (<i>In Press.</i>)	
Sudborough, J. J., and James, T. C. Practical Organic Chemistry.....	12mo,	*2 00
Suffling, E. R. Treatise on the Art of Glass Painting.....	8vo,	*3 50
Swan, K. Patents, Designs and Trade Marks. (Westminster Series.).....	8vo,	*2 00
Sweet, S. H. Special Report on Coal.....	8vo,	3 00
Swinburne, J., Wordingham, C. H., and Martin, T. C. Electric Currents. (Science Series No. 109.).....	16mo,	0 50
Swoope, C. W. Practical Lessons in Electricity.....	12mo,	*2 00
Tailfer, L. Bleaching Linen and Cotton Yarn and Fabrics. .	8vo,	*5 00
Tate, J. S. Surcharged and Different Forms of Retaining-walls. (Science Series No. 7.).....	16mo,	
Templeton, W. Practical Mechanic's Workshop Companion. .	12mo, mor.,	2 00
Terry, H. L. India Rubber and its Manufacture. (Westminster Series.).....	8vo,	*2 00
Thiess, J. B., and Joy, G. A. Toll Telephone Practice. (<i>In Preparation.</i>)		
Thom, C., and Jones, W. H. Telegraphic Connections. .	oblong 12mo,	1 50
Thomas, C. W. Paper-makers' Handbook.....	(<i>In Press.</i>)	
Thompson, A. B. Oil Fields of Russia.....	4to,	*7 50
— Petroleum Mining and Oil Field Development.	8vo,	*5 00
Thompson, E. P. How to Make Inventions.....	8vo,	0 50
Thompson, S. P. Dynamo Electric Machines. (Science Series No. 75.).....	16mo,	0 50
Thompson, W. P. Handbook of Patent Law of All Countries	16mo,	1 50
Thornley, T. Cotton Combing Machines.....	8vo,	*3 00
— Cotton Spinning	8vo,	
First Year.....		*1 50
Second Year.....		*2 50
Third Year.....		*2 50

34 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Thurso, J. W. Modern Turbine Practice.....	8vo,	*4 00
Tidy, C. Meymott. Treatment of Sewage. (Science Series No. 94.).....	16mo,	0 50
Tinney, W. H. Gold-mining Machinery.....	8vo,	*3 00
Titherley, A. W. Laboratory Course of Organic Chemistry.....	8vo,	*2 00
Toch, M. Chemistry and Technology of Mixed Paints.....	8vo,	*3 00
— Materials for Permanent Painting.....	12mo (<i>In Press.</i>)	
Todd, J., and Whall, W. B. Practical Seamanship.....	8vo,	*7 50
Tonge, J. Coal. (Westminster Series.).....	8vo,	*2 00
Townsend, F. Alternating Current Engineering....	8vo, boards,	*0 75
Townsend, J. Ionization of Gases by Collision.....	8vo,	*1 25
Transactions of the American Institute of Chemical Engineers.....	8vo,	
Vol. I. 1908.....		*6 00
Vol. II. 1909.....		*6 00
Traverse Tables. (Science Series No. 115.).....	16mo,	0 50
	mor.,	1 00
Trinks, W., and Housum, C. Shaft Governors. (Science Series No. 122.).....	16mo,	0 50
Trowbridge, W. P. Turbine Wheels. (Science Series No. 44.).....	16mo,	0 50
Tucker, J. H. A Manual of Sugar Analysis.....	8vo,	3 50
Tumlirz, O. Potential. Trans. by D. Robertson.....	12mo,	1 25
Tünner, P. A. Treatise on Roll-turning. Trans. by J. B. Pearse.....	8vo text and folio atlas,	10 00
Turbayne, A. A. Alphabets and Numerals.....	4to,	2 00
Turnbull, Jr., J., and Robinson, S. W. A Treatise on the Compound Steam-engine. (Science Series No. 8.).....	16mo,	
Turrill, S. M. Elementary Course in Perspective.....	12mo,	*1 25
Underhill, C. R. Solenoids, Electromagnets and Electromagnetic Windings.....	12mo,	*2 00
Urquhart, J. W. Electric Light Fitting.....	12mo,	2 00
— Electro-plating.....	12mo,	2 00
— Electrotyping.....	12mo,	2 00
— Electric Ship Lighting.....	12mo,	3 00
Universal Telegraph Cipher Code.....	12mo,	1 00

D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG 35

Vacher, F. Food Inspector's Handbook.....	12mo,	*2 50
Van Nostrand's Chemical Annual. Second issue 1909	12mo,	*2 50
— Year Book of Mechanical Engineering Data. First issue 1910.....	(In Press.)	
Van Wagenen, T. F. Manual of Hydraulic Mining.....	16mo,	1 00
Vega, Baron, Von Logarithmic Tables	8vo, half mor.,	2 50
Villon, A. M. Practical Treatise on the Leather Industry. Trans. by F. T. Addyman.....	8vo,	*10 00
Vincent, C. Ammonia and its Compounds. Trans. by M. J. Salter.....	8vo,	*2 00
Volk, C. Haulage and Winding Appliances.	8vo,	*4 00
Von Georgiowics, G. Chemical Technology of Textile Fibres.. Trans. by C. Salter.....	8vo,	*4 50
— Chemistry of Dyestuffs. Trans. by C. Salter.....	8vo,	*4 50
Vose, G. L. Graphic Method for Solving Certain Questions in Arithmetic and Algebra. (Science Series No. 16.)	16mo,	0 50
Wabner, R. Ventilation in Mines. Trans. by C. Salter. .	8vo,	*4 50
Wade, E. J. Secondary Batteries.....	8vo,	*4 00
Wadsworth, C. Primary Battery Ignition	12mo (In Press.)	
Wagner, E. Preserving Fruits, Vegetables, and Meat....	12mo,	*2 50
Walker, F. Aerial Navigation.....	8vo,	
— Dynamo Building. (Science Series No. 98.).....	16mo,	0 50
— Electric Lighting for Marine Engineers.....	8vo,	2 00
Walker, S. F. Steam Boilers, Engines and Turbines.....	8vo,	3 00
— Refrigeration, Heating and Ventilation on Shipboard.	12mo,	*2 00
— Electricity in Mining.....	8vo,	*3 50
Walker, W. H. Screw Propulsion.....	8vo,	0 75
Wallis-Taylor, A. J. Bearings and Lubrication.....	8vo,	*1 50
— Modern Cycles.....	8vo,	4 00
— Motor Cars.....	8vo,	1 80
— Motor Vehicles for Business Purposes.....	8vo,	3 50
— Pocket Book of Refrigeration and Ice Making.....	12mo,	1 50
— Refrigerating and Ice-making Machinery.....	8vo,	3 00
— Refrigeration and Cold Storage.....	8vo,	*4 50
— Sugar Machinery.....	12mo,	*2 00
Wanklyn, J. A. Treatise on the Examination of Milk ..	12mo,	1 00
— Water Analysis.....	12mo,	2 00

36 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Wansbrough, W. D. The A B C of the Differential Calculus	12mo,	*1 50
— Slide Valves	12mo,	*2 00
Ward, J. H. Steam for the Million	8vo,	1 00
Waring, Jr., G. E. Sanitary Conditions. (Science Series No. 31.)	16mo,	0 50
— Sewerage and Land Drainage		*6 00
— Modern Methods of Sewage Disposal	12mo,	2 00
— How to Drain a House	12mo,	1 25
Warren, F. D. Handbook on Reinforced Concrete	12mo,	*2 50
Watkins, A. Photography. (Westminster Series.)	8vo (<i>In Press.</i>)	
Watson, E. P. Small Engines and Boilers	12mo,	1 25
Watt, A. Electro-plating and Electro-refining of Metals		*4 50
— Electro-metallurgy	12mo,	1 00
— The Art of Paper Making		*3 00
— The Art of Soap-making	8vo,	3 00
— Leather Manufacture	8vo,	*4 00
Weale, J. Dictionary of Terms used in Architecture	12mo,	2 50
Weale's Scientific and Technical Series. (Complete list sent on application.)		
Weather and Weather Instruments	12mo,	1 00
	paper,	0 50
Webb, H. L. Guide to the Testing of Insulated Wires and Cables	12mo,	1 00
Webber, W. H. Y. Town Gas. (Westminster Series.)	8vo,	*2 00
Weekes, R. W. The Design of Alternate Current Transformers	12mo,	1 00
Weisbach, J. A Manual of Theoretical Mechanics	8vo,	*6 00
	sheep,	*7 50
Weisbach, J., and Herrmann, G. Mechanics of Air Machinery	8vo,	*3 75
Weston, E. B. Loss of Head Due to Friction of Water in Pipes	12mo,	*1 50
Weymouth, F. M. Drum Armatures and Commutators	8vo,	*3 00
Wheeler, J. B. Art of War	12mo,	1 75
— Field Fortifications	12mo,	1 75
Whipple, S. An Elementary and Practical Treatise on Bridge Building	8vo,	3 00

D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG 37

Whithard, P. Illuminating and Missal Painting	12mo,	1 50
Wilcox, R. M. Cantilever Bridges. (Science Series No. 25.)		
	16mo,	0 50
Wilkinson, H. D. Submarine Cable Laying and Repairing.		
	8vo,	*6 00
Williams, A. D., Jr., and Hutchinson, R. W. The Steam Turbine.		
	(In Press.)	
Williamson, R. S. On the Use of the Barometer	4to,	15 00
— Practical Tables in Meteorology and Hypsometry	4to,	2 50
Willson, F. N. Theoretical and Practical Graphics	4to,	*4 00
Wimperis, H. E. Internal Combustion Engine	8vo,	*3 00
Winchell, N. H., and A. N. Elements of Optical Mineralogy.		
	8vo,	*3 50
Winkler, C., and Lunge, G. Handbook of Technical Gas-Analy-		
sis	8vo,	4 00
Winslow, A. Stadia Surveying. (Science Series No. 77.).		
	16mo,	0 50
Wisser, Lieut. J. P. Explosive Materials. (Science Series No.		
70.)	16mo,	0 50
— Modern Gun Cotton. (Science Series No. 89.)	16mo,	0 50
Wood, De V. Luminiferous Aether. (Science Series No. 85.)		
	16mo,	0 50
Woodbury, D. V. Elements of Stability in the Well-propor-		
tioned Arch	8vo, half mor.,	4 00
Worden, E. C. The Nitrocellulose Industry. Two Volumes.		
	8vo (In Press.)	
Wright, A. C. Analysis of Oils and Allied Substances	8vo,	*3 50
— Simple Method for Testing Painter's Materials	8vo,	*2 50
Wright, H. E. Handy Book for Brewers	8vo,	*5 00
Wright, F. W. Design of a Condensing Plant	12mo,	*1 50
Wright, T. W. Elements of Mechanics	8vo,	*2 50
Wright, T. W., and Hayford, J. F. Adjustment of Observations		
	8vo,	*3 00
Young, J. E. Electrical Testing for Telegraph Engineers ...	8vo,	*4 00
Zahner, R. Transmission of Power. (Science Series No. 40.)		
	16mo,	
Zeidler, J., and Lustgarten, J. Electric Arc Lamps	8vo,	*2 00

38 D. VAN NOSTRAND COMPANY'S SHORT-TITLE CATALOG

Zeuner, A. Technical Thermodynamics. Trans. by J. F. Klein. Two Volumes.....	8vo,	*8 00
Zimmer, G. F. Mechanical Handling of Material.....	4to,	*10 00
Zipser, J. Textile Raw Materials. Trans. by C. Salter.....	8vo,	*5 00
Zur Nedden, F. Engineering Workshop Machines and Processes. Trans. by J. A. Davenport	8vo,	*2 00

UNIV. OF MICHIGAN,

MAY 22 1912

Books sent postpaid to any address on receipt of price.

Descriptive circulars and complete catalogs may be had on application.

FOSTER

ELECTRICAL

Completely Revised

ENGINEER'S

and Enlarged

POCKET BOOK

1800 pp. 1128 Ill. 718 Tables

Foster's Pocketbook is the busy engineer's first assistant. When he wants really reliable information, he turns to **Foster** to get it for him, and he gets it. **Foster** places before him data on the latest and best practice in electrical

engineering with a minimum expenditure of time and effort.

The material in **Foster** is logically arranged and is indexed in a full table of contents and a voluminous index that alone covers sixty pages, and to this is added a set of patent thumb index tabs that make reference to any section of the book practically instantaneous. The index is most thorough and reliable. It points right to the spot where the information sought is.

If you ever need information on electricity, you ought to have a copy of the new Fifth Edition, completely revised and enlarged, with *four-fifths* of the old matter replaced by new, up-to-date material, and containing all the excellent features named above, with 1,636 pages, 1,128 illustrations and 718 tables. The price is **\$5.00. ORDER A COPY NOW**

CIVIL ENGINEERS' POCKETBOOK

Nearly
Ready

By ALBERT I. FRYE, M. Am. Soc. C. E.

A COMPREHENSIVE treatment of Civil Engineering, in seventy sections, comprising about 1,400 pages; with 500 tables and 1,000 illustrations. Also a complete glossary of engineering terms.

Each main subject receives economic consideration and analysis, and is reinforced with excerpts from, and references to the most important cost and other data in our leading technical publications—including hundreds of illustrations of up-to-date engineering structures and details.

"How to design engineering structures economically" has received careful study in the preparation of this work. The fundamental principles laid down can be followed readily by every young engineer, and carried to practical completion in the finished structure.

Most of the tables are new and have been prepared regardless of time and expense. Many of them have been arranged in a novel manner which will appeal especially to engineers; and all have been carefully checked and rechecked to eliminate the possibility of errors.

Graphical methods have been illustrated freely throughout the work, in connection with analytical solutions.

Methods and cost of work have received special attention; and, thus, there is crowded into the 1,400 pages of text, in small type and in pocketbook form, a vast amount of engineering data, formulas and tables, that will be found absolutely indispensable to every engineer, contractor and student.

D. VAN NOSTRAND COMPANY, Publishers
23 Murray and 27 Warren Sts., **NEW YORK**

